

THE DYNKIN DIAGRAMS PACKAGE
VERSION 3.14159265

BEN MCKAY

CONTENTS

1. Quick introduction, 2
2. Set options globally, 3
3. Coxeter diagrams, 4
4. Satake diagrams, 5
5. How to fold, 6
6. Labels for the roots, 8
7. Label subscripts, 11
8. Height and depth of labels, 12
9. Text style for the labels, 12
10. Bracing roots, 13
11. Style, 17
12. Suppress or reverse arrows, 19
13. Backwards and upside down, 19
14. Drawing on top of a Dynkin diagram, 20
15. Mark lists, 21
16. Indefinite edges, 22
17. Parabolic subgroups, 24
18. Extended Dynkin diagrams, 26
19. Affine twisted and untwisted Dynkin diagrams, 27
20. Extended Coxeter diagrams, 28
21. Kac style, 29
22. Ceref style, 31
23. More on folded Dynkin diagrams, 32
24. Root ordering, 38
25. Connecting Dynkin diagrams, 38
26. Other examples, 41
27. Example: the complex simple Lie algebras, 49
28. An example of Mikhail Borovoi, 51
29. Syntax, 52
30. Options, 52
- References, 56

1. QUICK INTRODUCTION

Load the Dynkin diagram package (see options below)

```
\documentclass{amsart}
\usepackage{dynkin-diagrams}
\begin{document}
The Dynkin diagram of \(\mathbf{B}_3\) is \dynkin{B}{3}.
\end{document}
```

Invoke it

The Dynkin diagram of \(\mathbf{B}_3\) is \dynkin{B}{3}.

The Dynkin diagram of B_3 is $\bullet \rightarrowtail \bullet$.

Inside a *TikZ* statement

```
The Dynkin diagram of \(\mathbf{B}_3\) is
\tikz \dynkin{B}{3};
```

The Dynkin diagram of B_3 is $\bullet \rightarrowtail \bullet$

Inside a Dynkin diagram environment

```
The Dynkin diagram of \(\mathbf{B}_3\) is
\begin{dynkinDiagram}{B}{3}
\draw[very thick,red] (root 1) to [out=-45, in=-135] (root 3);
\end{dynkinDiagram}
```

The Dynkin diagram of B_3 is $\bullet \rightarrowtail \bullet$

Inside a *TikZ* environment

```
Baseline controls vertical alignment:
the Dynkin diagram of \(\mathbf{B}_3\) is
\begin{tikzpicture}[baseline=(origin.base)]
\dynkin{B}{3}
\draw[very thick,red] (root 1) to [out=-45, in=-135] (root 3);
\end{tikzpicture}
```

Baseline controls vertical alignment: the Dynkin diagram of B_3 is $\bullet \rightarrowtail \bullet$

Indefinite rank Dynkin diagrams

```
\dynkin{B}{}
```



Table 1: The Dynkin diagrams of the reduced simple root systems [3] pp. 265–290, plates I–IX

A_n		<code>\dynkin{A}{}</code>
C_n		<code>\dynkin{C}{}</code>
D_n		<code>\dynkin{D}{}</code>
E_6		<code>\dynkin{E}{6}</code>
E_7		<code>\dynkin{E}{7}</code>
E_8		<code>\dynkin{E}{8}</code>
F_4		<code>\dynkin{F}{4}</code>
G_2		<code>\dynkin{G}{2}</code>

2. SET OPTIONS GLOBALLY

Most options set globally ...

```
\pgfkeys{/Dynkin diagram,edge length=.5cm,fold radius=.5cm,
indefinite edge/.style={%
    draw=black,fill=white,thin,densely dashed}}
```

You can also pass options to the package in `\usepackage`. *Danger:* spaces in option names are replaced with hyphens: `edge length=1cm` is `edge-length=1cm` as a global option; moreover you should drop the extension .style on any option with spaces in its name (but not otherwise). For example,


... or pass global options to the package

```
\usepackage[%
    ordering=Kac,
    edge/.style=blue,
    indefinite-edge={draw=green,fill=white,densely dashed},
    indefinite-edge-ratio=5,
    mark=o,
    root-radius=.06cm]
{dynkin-diagrams}
```

3. COXETER DIAGRAMS

Coxeter diagram option

\dynkin[Coxeter]{F}{4}

gonality option for G_2 and I_n Coxeter diagrams

\begin{aligned} \text{\(G_2=\)}\dynkin[Coxeter,gonality=n]{G}{2}\text{\(), } \\ \text{\(I_n=\)}\dynkin[Coxeter,gonality=n]{I}{}\text{\)}\end{aligned}

$$G_2 = \bullet^n \bullet, \quad I_n = \bullet^n \bullet$$

Table 2: The Coxeter diagrams of the simple reflection groups

A_n		\dynkin[Coxeter]{A}{}
B_n		\dynkin[Coxeter]{B}{}
C_n		\dynkin[Coxeter]{C}{}
E_6		\dynkin[Coxeter]{E}{6}
E_7		\dynkin[Coxeter]{E}{7}
E_8		\dynkin[Coxeter]{E}{8}
F_4		\dynkin[Coxeter]{F}{4}
G_2		\dynkin[Coxeter,gonality=n]{G}{2}
H_3		\dynkin[Coxeter]{H}{3}
H_4		\dynkin[Coxeter]{H}{4}
I_n		\dynkin[Coxeter,gonality=n]{I}{}

4. SATAKE DIAGRAMS

Satake diagrams use the standard name instead of a rank

```
\(A_{IIIb}=\dynkin{A}{IIIb}\)
```

$$A_{IIIb} = \begin{array}{c} \square \\ \square \end{array}$$






We use a solid gray bar to denote the folding of a Dynkin diagram, rather than the usual double arrow, since the diagrams turn out simpler and easier to read.

Table 3: The Satake diagrams of the real simple Lie algebras [13] p. 532–534

A_I		<code>\dynkin{A}{I}</code>
A_{II}		<code>\dynkin{A}{II}</code>
A_{IIIa}		<code>\dynkin{A}{IIIa}</code>
A_{IIIb}		<code>\dynkin{A}{IIIb}</code>
A_{IV}		<code>\dynkin{A}{IV}</code>
B_I		<code>\dynkin{B}{I}</code>
B_{II}		<code>\dynkin{B}{II}</code>
C_I		<code>\dynkin{C}{I}</code>
C_{IIa}		<code>\dynkin{C}{IIa}</code>
C_{IIb}		<code>\dynkin{C}{IIb}</code>
D_{Ia}		<code>\dynkin{D}{Ia}</code>
D_{Ib}		<code>\dynkin{D}{Ib}</code>
D_{Ic}		<code>\dynkin{D}{Ic}</code>
D_{II}		<code>\dynkin{D}{II}</code>
D_{IIIa}		<code>\dynkin{D}{IIIa}</code>
D_{IIIb}		<code>\dynkin{D}{IIIb}</code>
E_I		<code>\dynkin{E}{I}</code>
E_{II}		<code>\dynkin{E}{II}</code>

continued ...


Table 3: ... continued

E_{III}		\dynkin{E}{III}
E_{IV}		\dynkin{E}{IV}
E_V		\dynkin{E}{V}
E_{VI}		\dynkin{E}{VI}
E_{VII}		\dynkin{E}{VII}
E_{VIII}		\dynkin{E}{VIII}
E_{IX}		\dynkin{E}{IX}
F_I		\dynkin{F}{I}
F_{II}		\dynkin{F}{II}
G_I		\dynkin{G}{I}

5. HOW TO FOLD


If you don't like the solid gray "folding bar", most people use arrows. Here is E_{II}

```
\newcommand{\invol}[2]{\draw[latex-latex] (root #1) to
 [out=-60,in=-120] node[midway,below]{$\sigma$} (root #2);}
\begin{dynkinDiagram}[edge length=.75cm,labels*={1,...,6}]{E}{6}
\invol{1}{6}\invol{3}{5}
\end{dynkinDiagram}
```




The double arrows for A_{IIIa} are big

```
\newcommand{\invol}[2]{\draw[latex-latex] (root #1) to
 [out=-60,in=-120] node[midway,below]{$\sigma$} (root #2);}
\begin{dynkinDiagram}[edge length=.75cm]{A}{oo.o**.*o.oo}
\invol{1}{10}\invol{2}{9}\invol{3}{8}\invol{4}{7}\invol{5}{6}
\end{dynkinDiagram}
```




If you don't like the solid gray "folding bar", most people use arrows ...

```
\tikzset{/Dynkin diagram/fold style/.style={stealth-stealth,thick,
shorten <=1mm,shorten >=1mm,}}
\dynkin[ply=3,edge length=.75cm]{D}{4}
\begin{dynkinDiagram}[ply=4]{D}[1]%
{****.*****.*****}
\dynkinFold{1}{13}
\dynkinFold[bend right=90]{0}{14}
\end{dynkinDiagram}
```



...but you could try springs pulling roots together


```
\tikzset{/Dynkin diagram/fold style/.style=
{decorate,decoration={name=coil,aspect=0.5,
segment length=1mm,amplitude=.6mm}}}
\dynkin[ply=3,edge length=.75cm]{D}{4}
\begin{dynkinDiagram}[ply=4]{D}[1]%
{****,*****,*}
\dynkinFold{1}{13}
\dynkinFold[bend right=90]{0}{14}
\end{dynkinDiagram}
```



6. LABELS FOR THE ROOTS


Make a macro to assign labels to roots

```
\dynkin[label,label macro/.code={\alpha_{{\mathrlap{\#1}}}},edge
length=.75cm]{D}{5}
```



Labelling several roots

```
\dynkin[labels={2,...,5,,7},label
macro/.code={\alpha_{{\mathrlap{\#1}}}}]{A}{7}
```




The `foreach` notation I

```
\dynkin[labels={1,3,...,7},]{A}{9}
```




The `foreach` notation II

```
\dynkin[labels={,\alpha_2,\alpha_3,\alpha_4,\alpha_5,\alpha_6,\alpha_7},]{A}{7}
```

The `foreach` notation III

```
\dynkin[label
macro/.code={\beta_{\mathrlap{\#1}}},labels={,2,...,7},]{A}{7}
```



Label the roots individually by root number

```
\dynkin[label]{B}{3}
```



Label a single root

```
\begin{dynkinDiagram}{B}{3}
\dynkinLabelRoot{2}{\alpha_2}
\end{dynkinDiagram}
```




Access root labels via TikZ

```
\begin{dynkinDiagram}{B}{3}
\node[below] at (root 2) {\alpha_2};
\end{dynkinDiagram}
```




Commands to label several roots

```
\begin{dynkinDiagram}{A}{7}
\dynkinLabelRoots{\alpha_2,\alpha_3,\alpha_4,\alpha_5,,\alpha_7}
\end{dynkinDiagram}
```




The labels have default locations, mostly below roots

```
\dynkin[edge length=.75cm,labels={1,2,3}]{E}{8}
```




The starred form flips labels to alternate locations, mostly above roots

```
\dynkin[edge length=.75cm,labels*={1,2,3}]{E}{8}
```




Labelling several roots and alternates

```
\dynkin[%
label macro/.code={\alpha_{\mathrlap{\#1}}},
label macro*/.code={\gamma_{\mathrlap{\#1}}},
labels={,2,...,5,,7},
labels*={1,3,4,5,6}]{A}{7}
```



Commands to label several roots

```
\begin{dynkinDiagram}{A}{7}
\dynkinLabelRoots{\alpha_2,\alpha_3,\alpha_4,\alpha_5,,\alpha_7}
\dynkinLabelRoots*{a,b,c,d,e,f,g}
\end{dynkinDiagram}
```




7. LABEL SUBSCRIPTS

Note the slight improvement that `\mathrlap` makes: the labels are centered on the middle of the letter α , ignoring the space taken up by the subscripts.

Label spacing improvement

```
\dynkin[label,label macro/.code={\alpha_{#1}},edge length=.75cm]{D}{15}
\par\noindent{}%
\dynkin[label,label macro/.code={\alpha_{\mathrlap{#1}}},edge
length=.75cm]{D}{15}
```




8. HEIGHT AND DEPTH OF LABELS

Labels are set with default maximum height the height of the character b , and default maximum depth the depth of the character g . To change these, set `label height` and `label depth`:

Change height and dept of characters

```
\dynkin[labels={a,b,c,d}]{F}{4}
\dynkin[labels*={a,b,c,d}]{F}{4}
\dynkin[%]
label macro/.code={\alpha_{{\mathrlap{\#1}}}},
label macro*/.code={\gamma_{{\mathrlap{\#1}}}},
label height=$\alpha_1$,
label depth=$\alpha_1$,
labels={,2,...,5,,7},
labels*={1,3,4,5,6}{A}{7}
\dynkin[labels={A,B,C,D},label height=$A$,label depth=$A$]{F}{4}
\dynkin[labels={a^1,b^2,c^3,d^4},label height=$X^X$]{F}{4}
```



9. TEXT STYLE FOR THE LABELS

Use a text style: big and blue

```
\begin{dynkinDiagram}[text style={scale=1.2,blue},
edge length=.75cm,
labels={1,2,n-1,n},
label macro/.code={\alpha_{{\mathrlap{\#1}}}}
]{A}{}
\end{dynkinDiagram}
```



Use a text style; font selection is in the label macro


```
\begin{dynkinDiagram}[text style={scale=1.2,blue},
edge length=.75cm,
labels={1,2,n-1,n},
label macro/.code={\mathbb{A}_{{\mathrlap{\#1}}}}]{A}{}
\end{dynkinDiagram}
```



10. BRACING ROOTS


Bracing roots

```
\begin{dynkinDiagram}{A}{*,*x*.*}
\dynkinBrace[p]{1}{2}
\dynkinBrace[q]{4}{5}
\end{dynkinDiagram}
```



Bracing roots, and a starred form

```
\begin{dynkinDiagram}{A}{10}
\dynkinBrace[\text{Roots 2 to 9}]{2}{9}
\dynkinBrace*[\text{Roots 3 to 8}]{3}{8}
\end{dynkinDiagram}
```



Bracing roots

```
\newcommand\circleRoot[1]{\draw (root #1) circle (3pt);}
\begin{dynkinDiagram}{A}{**.***.***.***.***.*}
\circleRoot{4}\circleRoot{7}\circleRoot{10}\circleRoot{13}
\dynkinBrace[y-1]{1}{3}
\dynkinBrace[z-1]{5}{6}
\dynkinBrace[t-1]{11}{12}
\dynkinBrace[x-1]{14}{16}
\end{dynkinDiagram}
```






Table 4: Dynkin diagrams from Euler products [17]




continued . . .

Table 4: ... continued




continued ...

Table 4: ... continued



continued ...

Table 4: ... continued



```
\tikzset{/Dynkin diagram,ordering=Dynkin,label macro/.code={\alpha_{##1}}}
\newcounter{EPNo}
\setcounter{EPNo}{0}
\NewDocumentCommand{\EP}{smmmmm}%
{%
\stepcounter{EPNo}\roman{EPNo}. \&
\def\eL{.6cm}
\IfStrEqCase{#2}{%
{D}{\gdef\eL{1cm}}%
{E}{\gdef\eL{.75cm}}%
{F}{\gdef\eL{.35cm}}%
{G}{\gdef\eL{.35cm}}%
}%
\tikzset{/Dynkin diagram,edge length=\eL}
\IfBooleanTF{#1}%
{\dynkin[backwards,labels*={#4},labels={#5}]{#2}{#3}}%
{\dynkin[labels*={#4},labels={#5}]{#2}{#3}}%
\\
}%
\begin{longtable}{MM}
\caption{Dynkin diagrams from Euler products \cite{Langlands:1967}}\\
\endfirsthead
\caption{\dots continued}\\
\endhead
\multicolumn{2}{c}{continued \dots}\\
\endfoot
\endlastfoot
\EP{A}{***.**}{1,1,1,1,1}{1,2,n-1,n}
\EP{A}{***.**}{1,1,1,1,1}{1,2,n-1,n}
\EP{A}{**.***.*}{1,1,1,1,1,1}{1,2,m-1,,m,n}
\EP{B}{**.****}{2,2,2,2,1}{1,2,n-1,n}
\EP*{B}{***.**}{2,2,2,2,1}{n,n-1,2,1,}
\EP{C}{**.****}{1,1,1,1,2}{1,2,n-1,}
\EP*{C}{***.**}{1,1,1,1,2}{n,n-1,2,1,}
\EP{D}{**.*****}{1,1,1,1,1,1}{1,2,n-2,n-1,n}
\EP{D}{**.*****}{1,1,1,1,1,1}{1,2,n-2,n-1,n}
```

```
\EP{E}{6}{1,1,1,1,1,1}{1,...,5}
\EP*{E}{7}{1,1,1,1,1,1}{6,...,1}
\EP{E}{7}{1,1,1,1,1,1}{1,...,6}
\EP*{E}{8}{1,1,1,1,1,1,1}{7,...,1}
\EP{E}{8}{1,1,1,1,1,1,1}{1,...,7}
\EP{G}{2}{1,3}{,1}
\EP{G}{2}{1,3}{1}
\EP{B}{**.***}{2,2,2,2,1}{1,2,n-1,n}
\EP{F}{4}{1,1,2,2}{3,2,1}
\EP{C}{3}{1,1,2}{,2,1}
\EP{C}{**.***}{1,1,1,1,2}{1,n-2,n-1,n}
\EP*{B}{3}{2,2,1}{1,2}
\EP{F}{4}{1,1,2,2}{1,2,3}
\EP{D}{**.****}{1,1,1,1,1,1}{1,2,n-2,n-2,n,n}
\EP{E}{6}{1,1,1,1,1,1}{1,2,3,4,,5}
\EP{E}{6}{1,1,1,1,1,1}{1,2,3,5,,4}
\EP*{E}{7}{1,1,1,1,1,1}{5,...,1,6}
\EP*{E}{7}{1,1,1,1,1,1}{6,4,3,2,1,5}
\EP*{E}{8}{1,1,1,1,1,1,1}{6,...,1,7}
\EP*{E}{8}{1,1,1,1,1,1,1}{7,5,4,3,2,1,6}
\EP*{E}{7}{1,1,1,1,1,1,1}{5,...,1,,6}
\EP*{E}{7}{1,1,1,1,1,1,1}{1,...,5,,6}
\EP*{E}{8}{1,1,1,1,1,1,1}{6,...,1,,7}
\end{longtable}
```

11. STYLE

Colours

```
\dynkin[
    edge/.style={blue!50,thick},
    */.style=blue!50!red,
    arrow color=red]{F}{4}
```




Edge lengths

The Dynkin diagram of $\backslash(A_3\backslash)$ is \dynkin[edge
length=1.2,parabolic=3]{A}{3}

The Dynkin diagram of A_3 is

Root marks

```
\dynkin{E}{8}
\dynkin[mark=*]{E}{8}
\dynkin[mark=o]{E}{8}
\dynkin[mark=O]{E}{8}
\dynkin[mark=t]{E}{8}
\dynkin[mark=x]{E}{8}
\dynkin[mark=X]{E}{8}
```



At the moment, you can only use:

- * solid dot
- o hollow circle
- O double hollow circle
- t tensor root
- x crossed root
- X thickly crossed root

Mark styles

The parabolic subgroup $\langle E_{8,124} \rangle$ is
 $\backslash\text{dynkin}[parabolic=124,x/.style=\{brown,very thick\}]{E}{8}$

The parabolic subgroup $E_{8,124}$ is 

Sizes of root marks

```
\langle A_{3,3} \rangle with big root marks is \dynkin[root
radius=.08cm,parabolic=3]{A}{3}
```

$A_{3,3}$ with big root marks is 

12. SUPPRESS OR REVERSE ARROWS

Some diagrams have double or triple edges

```
\dynkin{F}{4}
\dynkin{G}{2}
```



Suppress arrows

```
\dynkin[arrows=false]{F}{4}
\dynkin[arrows=false]{G}{2}
```



Reverse arrows

```
\dynkin[reverse arrows]{F}{4}
\dynkin[reverse arrows]{G}{2}
```



13. BACKWARDS AND UPSIDE DOWN

Default

```
\dynkin{E}{8}
\dynkin{F}{4}
\dynkin{G}{2}
```



Backwards

```
\dynkin[backwards]{E}{8}
\dynkin[backwards]{F}{4}
\dynkin[backwards]{G}{2}
```



Reverse arrows

```
\dynkin[reverse arrows]{F}{4}
\dynkin[reverse arrows]{G}{2}
```




Backwards, reverse arrows

```
\dynkin[backwards,reverse arrows]{F}{4}
\dynkin[backwards,reverse arrows]{G}{2}
```



Backwards versus upside down

```
\dynkin[label]{E}{8}
\dynkin[label,backwards]{E}{8}
\dynkin[label,upside down]{E}{8}
\dynkin[label,backwards,upside down]{E}{8}
```



14. DRAWING ON TOP OF A DYNKIN DIAGRAM


TikZ can access the roots themselves

```
\begin{dynkinDiagram}{A}{4}
\fill[white,draw=black] (root 2) circle (.15cm);
\fill[white,draw=black] (root 2) circle (.1cm);
\draw[black] (root 2) circle (.05cm);
\end{dynkinDiagram}
```




Draw curves between the roots

```
\begin{dynkinDiagram}[label]{E}{8}
    \draw[very thick, black!50,-latex]
        (root 3.south) to [out=-45, in=-135] (root 6.south);
\end{dynkinDiagram}
```



Change marks

```
\begin{dynkinDiagram}[mark=o,label]{E}{8}
    \dynkinRootMark{*}{5}
    \dynkinRootMark{*}{8}
\end{dynkinDiagram}
```



15. MARK LISTS

The package allows a list of root marks instead of a rank:

A mark list

```
\dynkin{E}{oo**ttxx}
```



The mark list `oo**ttxx` has one mark for each root: o, o, ..., x. Roots are listed in the current default ordering. (Careful: in an affine root system, a mark list will *not* contain a mark for root zero.)

If you need to repeat a mark, you can give a *single digit* positive integer to indicate how many times to repeat it.

A mark list with repetitions

```
\dynkin{A}{x4o3t4}
```




Table 5: Classical Lie superalgebras [10]. We need a slightly larger root radius parameter to distinguish the tensor product symbols from the solid dots.

	\tikzset{/Dynkin diagram,root radius=.07cm}
A_{mn}	$\circ\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\circ$ \dynkin{A}{o3.oto.oo}
B_{mn}	$\circ\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\otimes\circ$ \dynkin{B}{o3.oto.oo}
B_{0n}	$\circ\cdots\circ\circ\cdots\circ\circ\cdots\circ\otimes\bullet$ \dynkin{B}{o3.o3.o*}
C_n	$\otimes\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\otimes\circ$ \dynkin{C}{too.oto.oo}
D_{mn}	$\circ\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\circ\circ\circ$ \dynkin{D}{o3.oto.o4}
$D_{21\alpha}$	$\circ\otimes\circ$ \dynkin{A}{oto}
F_4	$\circ\circ\otimes\circ\otimes$ \dynkin{F}{ooot}
G_3	$\otimes\bullet\bullet$ \dynkin[extended,affine mark=t, reverse arrows]{G}{2}

Table 6: Classical Lie superalgebras [10]. Here we see the problem with using the default root radius parameter, which is too small for tensor product symbols.

A_{mn}	$\circ\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\circ$	\dynkin{A}{o3.oto.oo}
B_{mn}	$\circ\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\otimes\circ$	\dynkin{B}{o3.oto.oo}
B_{0n}	$\circ\cdots\circ\circ\cdots\circ\circ\cdots\circ\otimes\bullet$	\dynkin{B}{o3.o3.o*}
C_n	$\otimes\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\otimes\circ$	\dynkin{C}{too.oto.oo}
D_{mn}	$\circ\cdots\circ\circ\cdots\circ\otimes\circ\cdots\circ\circ\circ\circ$	\dynkin{D}{o3.oto.o4}
$D_{21\alpha}$	$\circ\otimes\circ$	\dynkin{A}{oto}
F_4	$\circ\circ\otimes\circ\otimes$	\dynkin{F}{ooot}
G_3	$\otimes\bullet\bullet$	\dynkin[extended,affine mark=t, reverse arrows]{G}{2}

16. INDEFINITE EDGES

An *indefinite edge* is a dashed edge between two roots, $\bullet\cdots\bullet$ indicating that an indefinite number of roots have been omitted from the Dynkin diagram. In between any two entries in a mark list, place a period to indicate an indefinite edge:

Indefinite edges


```
\dynkin{D}{o.o*.*.t.to.t}
```

The diagram shows a Dynkin type D Dynkin diagram. It consists of five nodes connected by solid lines. Between the 4th and 5th nodes, there is a dashed line with a dot at each end, representing an indefinite edge. The nodes are labeled with circles: open circles for simple roots and a solid circle for the long root.

In certain diagrams, roots may have an edge between them even though they are not subsequent in the ordering. For such rare situations, there is an option:


Indefinite edge option

```
\dynkin[make indefinite edge={3-5},label]{D}{5}
```




Give a list of edges to become indefinite

```
\dynkin[make indefinite edge/.list={1-2,3-5},label]{D}{5}
```



Indefinite edge style

```
\dynkin[indefinite edge/.style={draw=black,fill=white,thin,densely dashed},%
edge length=1cm,%
make indefinite edge={3-5}%
{D}{5}
```



The ratio of the lengths of indefinite edges to those of other edges

```
\dynkin[edge length = .5cm,%
indefinite edge ratio=3,%
make indefinite edge={3-5}%
{D}{5}
```




Table 7: Springer's table of indices [24], pp. 320-321, with one form of E_7 corrected

A_n		
A_n		
B_n		
C_n		
D_n		
E_6		\dynkin{E}{*oooo*}
E_6		\dynkin{E}{o*o*oo}
E_6		\dynkin{E}{o*oooo}
E_6		\dynkin{E}{**oooo*}
E_7		\dynkin{E}{*oooooooo}
E_7		\dynkin{E}{oooooo*o}
E_7		\dynkin{E}{ooooooo*}
E_7		\dynkin{E}{*oooooo*o}
E_7		\dynkin{E}{*ooooo**}
E_7		\dynkin{E}{*oooo*o}
E_8		\dynkin{E}{*oooooooo}
E_8		\dynkin{E}{oooooooo*}
E_8		\dynkin{E}{*ooooooo*}
E_8		\dynkin{E}{ooooooo**}
E_8		\dynkin{E}{*ooooo***}
F_4		\dynkin{F}{ooo*}
D_4		\dynkin{D}{o*oo}


17. PARABOLIC SUBGROUPS

Each set of roots is assigned a number, with each binary digit zero or one to say whether the corresponding root is crossed or not:

The flag variety of pointed lines in
projective 3-space is associated to
the Dynkin diagram `\dynkin[parabolic=3]{A}{3}`.

The flag variety of pointed lines in projective 3-space is associated to the
Dynkin diagram $\text{---} \times \text{---} \bullet$.

Table 8: The Hermitian symmetric spaces

A_n		Grassmannian of k -planes in \mathbb{C}^{n+1}
B_n		$(2n-1)$ -dimensional hyperquadric, i.e. the variety of null lines in \mathbb{C}^{2n+1}
C_n		space of Lagrangian n -planes in \mathbb{C}^{2n}
D_n		$(2n-2)$ -dimensional hyperquadric, i.e. the variety of null lines in \mathbb{C}^{2n}
D_n		one component of the variety of maximal dimension null subspaces of \mathbb{C}^{2n}
D_n		the other component
E_6		complexified octave projective plane
E_6		its dual plane
E_7		the space of null octave 3-planes in octave 6-space

```
\NewDocumentCommand{\HSS}{m}{\IfNoValueTF{#2}{\dynkin{#3}{#4}}{\dynkin[parabolic=#2]{#3}{#4}}\And \\}
\renewcommand*{\arraystretch}{1.5}
\begin{longtable}{|c|c|c|} \caption{The Hermitian symmetric spaces} \endfirsthead
\caption{dots continued}\And \endhead
\caption{continued dots}\And \endfoot
\endlastfoot
\HSS{A_n}[A]{**.*x*.*}{Grassmannian of $k$-planes in $\mathbb{C}^{n+1}$}
\HSS{B_n}[1][B]{$(2n-1)$-dimensional hyperquadric, i.e. the variety of null lines in $\mathbb{C}^{2n+1}$}
\HSS{C_n}[16][C]{space of Lagrangian $n$-planes in $\mathbb{C}^{2n}$}
\HSS{D_n}[1][D]{$(2n-2)$-dimensional hyperquadric, i.e. the variety of null lines in $\mathbb{C}^{2n}$}
\HSS{D_n}[32][D]{one component of the variety of maximal dimension null subspaces of $\mathbb{C}^{2n}$}
\HSS{D_n}[16][D]{the other component}
\HSS{E_6}[1][E]{complexified octave projective plane}
\HSS{E_6}[32][E]{its dual plane}
\HSS{E_7}[64][E]{the space of null octave 3-planes in octave 6-space}
\end{longtable}
```

Folded parabolics look bad (zoom in on a root)

```
\dynkin[fold,parabolic=3]{C}{2}
\dynkin[fold,parabolic=3]{G}{2}
```



Folded parabolics: you can try using thicker crosses


```
\dynkin[fold,x/.style={very thick,line cap=round},parabolic=3]{C}{2}
\dynkin[fold,x/.style={ultra thick,line cap=round},parabolic=3]{G}{2}
```



18. EXTENDED DYNKIN DIAGRAMS

Extended Dynkin diagrams

```
\dynkin[extended]{A}{7}
```



The extended Dynkin diagrams are also described in the notation of Kac [15] p. 55 as affine untwisted Dynkin diagrams: we extend `\dynkin{A}{7}` to become `\dynkin{A}{1}{7}`:

Extended Dynkin diagrams

```
\dynkin{A}{1}{7}
```




Table 9: The Dynkin diagrams of the extended simple root systems

A_1^1	$\Leftrightarrow \bullet$	<code>\dynkin[extended]{A}{1}</code>
A_n^1		<code>\dynkin[extended]{A}{}</code>
B_n^1		<code>\dynkin[extended]{B}{}</code>

continued ...

Table 9: ... continued

C_n^1		<code>\dynkin[extended]{C}{}</code>
D_n^1		<code>\dynkin[extended]{D}{}</code>
E_6^1		<code>\dynkin[extended]{E}{6}</code>
E_7^1		<code>\dynkin[extended]{E}{7}</code>
E_8^1		<code>\dynkin[extended]{E}{8}</code>
F_4^1		<code>\dynkin[extended]{F}{4}</code>
G_2^1		<code>\dynkin[extended]{G}{2}</code>

19. AFFINE TWISTED AND UNTWISTED DYNKIN DIAGRAMS

The affine Dynkin diagrams are described in the notation of Kac [15] p. 55:

Affine Dynkin diagrams

```
\(A^{(1)}_7=\dynkin{A}[1]{7}, \
E^{(2)}_6=\dynkin{E}[2]{6}, \
D^{(3)}_4=\dynkin{D}[3]{4})\)
```

$$A_7^{(1)} = \text{Diagram of } A_7^{(1)}, \quad E_6^{(2)} = \text{Diagram of } E_6^{(2)}, \quad D_4^{(3)} = \text{Diagram of } D_4^{(3)}$$

Table 10: The affine Dynkin diagrams

A_1^1		<code>\dynkin{A}[1]{1}</code>
A_n^1		<code>\dynkin{A}[1]{}</code>
B_n^1		<code>\dynkin{B}[1]{}</code>
C_n^1		<code>\dynkin{C}[1]{}</code>
D_n^1		<code>\dynkin{D}[1]{}</code>
E_6^1		<code>\dynkin{E}[1]{6}</code>
E_7^1		<code>\dynkin{E}[1]{7}</code>
E_8^1		<code>\dynkin{E}[1]{8}</code>
F_4^1		<code>\dynkin{F}[1]{4}</code>

continued ...

Table 12: The extended (affine) Coxeter diagrams

A_n		<code>\dynkin[extended,Coxeter]{A}{}{}</code>
B_n		<code>\dynkin[extended,Coxeter]{B}{}{}</code>
C_n		<code>\dynkin[extended,Coxeter]{C}{}{}</code>
D_n		<code>\dynkin[extended,Coxeter]{D}{}{}</code>
E_6		<code>\dynkin[extended,Coxeter]{E}{6}{}</code>
E_7		<code>\dynkin[extended,Coxeter]{E}{7}{}</code>
E_8		<code>\dynkin[extended,Coxeter]{E}{8}{}</code>
F_4		<code>\dynkin[extended,Coxeter]{F}{4}{}</code>
G_2		<code>\dynkin[extended,Coxeter]{G}{2}{}</code>
H_3		<code>\dynkin[extended,Coxeter]{H}{3}{}</code>
H_4		<code>\dynkin[extended,Coxeter]{H}{4}{}</code>
I_1		<code>\dynkin[extended,Coxeter]{I}{1}{}</code>

21. KAC STYLE

We include a style called **Kac** which tries to imitate the style of [15].

Kac style

```
\dynkin[Kac]{F}{4}
```

Table 13: The Dynkin diagrams of the simple root systems in Kac style

A_n		<code>\dynkin[A]{}{}</code>
B_n		<code>\dynkin[B]{}{}</code>
C_n		<code>\dynkin[C]{}{}</code>
D_n		<code>\dynkin[D]{}{}</code>

continued ...

Table 13: ... continued

E_6		<code>\dynkin{E}{6}</code>
E_7		<code>\dynkin{E}{7}</code>
E_8		<code>\dynkin{E}{8}</code>
F_4		<code>\dynkin{F}{4}</code>
G_2		<code>\dynkin{G}{2}</code>

Table 14: The Dynkin diagrams of the extended simple root systems in Kac style

A_1^1		<code>\dynkin[extended]{A}{1}</code>
A_n^1		<code>\dynkin[extended]{A}{}{}</code>
B_n^1		<code>\dynkin[extended]{B}{}{}</code>
C_n^1		<code>\dynkin[extended]{C}{}{}</code>
D_n^1		<code>\dynkin[extended]{D}{}{}</code>
E_6^1		<code>\dynkin[extended]{E}{6}</code>
E_7^1		<code>\dynkin[extended]{E}{7}</code>
E_8^1		<code>\dynkin[extended]{E}{8}</code>
F_4^1		<code>\dynkin[extended]{F}{4}</code>
G_2^1		<code>\dynkin[extended]{G}{2}</code>

Table 15: The Dynkin diagrams of the twisted simple root systems in Kac style

A_2^2	$\circ \Leftrightarrow \circ$	<code>\dynkin[A]{2}{2}</code>
A_{ev}^2	$\circ \Leftarrow \circ - \circ - \circ - \cdots - \circ - \circ \Leftarrow \circ$	<code>\dynkin[A]{2}{even}</code>
A_{od}^2		<code>\dynkin[A]{2}{odd}</code>
D_n^2	$\circ \Leftarrow \circ - \circ - \cdots - \circ - \circ \Rightarrow \circ$	<code>\dynkin[D]{2}{}</code>
E_6^2	$\circ - \circ - \circ \Leftarrow \circ - \circ$	<code>\dynkin[E]{2}{6}</code>
D_4^3	$\circ - \circ \Leftrightarrow \circ$	<code>\dynkin[D]{3}{4}</code>

22. CEREF STYLE

We include a style called `ceref` which paints oblong root markers with shadows. The word “ceref” is an old form of the word “serif”.

Ceref style

```
\dynkin[ceref]{F}{4}
```



Table 16: The Dynkin diagrams of the simple root systems in ceref style






A_n		<code>\dynkin[A]{}{}</code>
B_n		<code>\dynkin[B]{}{}</code>
C_n		<code>\dynkin[C]{}{}</code>
D_n		<code>\dynkin[D]{}{}</code>
E_6		<code>\dynkin[E]{6}{}</code>
E_7		<code>\dynkin[E]{7}{}</code>
E_8		<code>\dynkin[E]{8}{}</code>
F_4		<code>\dynkin[F]{4}{}</code>
G_2		<code>\dynkin[G]{2}{}</code>

Table 17: The Dynkin diagrams of the extended simple root systems in ceref style

A_1^1		<code>\dynkin[extended]{A}{1}</code>
A_n^1		<code>\dynkin[extended]{A}{}{}</code>
B_n^1		<code>\dynkin[extended]{B}{}{}</code>
C_n^1		<code>\dynkin[extended]{C}{}{}</code>
D_n^1		<code>\dynkin[extended]{D}{}{}</code>
E_6^1		<code>\dynkin[extended]{E}{6}</code>
E_7^1		<code>\dynkin[extended]{E}{7}</code>
E_8^1		<code>\dynkin[extended]{E}{8}</code>
F_4^1		<code>\dynkin[extended]{F}{4}</code>
G_2^1		<code>\dynkin[extended]{G}{2}</code>

Table 18: The Dynkin diagrams of the twisted simple root systems in ceref style


A_2^2		<code>\dynkin{A}{2}{2}</code>
A_{ev}^2		<code>\dynkin{A}{2}{even}</code>
A_{od}^2		<code>\dynkin{A}{2}{odd}</code>
D_n^2		<code>\dynkin{D}{2}{}{}</code>
E_6^2		<code>\dynkin{E}{2}{6}</code>
D_4^3		<code>\dynkin{D}{3}{4}</code>

23. MORE ON FOLDED DYNKIN DIAGRAMS

The Dynkin diagrams package has limited support for folding Dynkin diagrams.


Folding

```
\dynkin{fold}{A}{13}
```




Big fold radius

```
\dynkin[fold,fold radius=1cm]{A}{13}
```



Small fold radius

```
\dynkin[fold,fold radius=.2cm]{A}{13}
```



Some Dynkin diagrams have multiple foldings, which we attempt to distinguish (not entirely successfully) by their *ply*: the maximum number of roots folded together. Most diagrams can only allow a 2-ply folding, so `fold` is a synonym for `ply=2`.

3-ply

```
\dynkin[ply=3]{D}{4}
\dynkin[ply=3,fold right]{D}{4}
\dynkin[ply=3]{D}{1}{4}
```



4-ply


```
\dynkin[ply=4]{D}{1}{4}
```



The $D_\ell^{(1)}$ diagrams can be folded on their left end and separately on their right end:

Left, right and both

```
\dynkin{D}[1]{} \
\dynkin[fold left]{D}[1]{} \
\dynkin[fold right]{D}[1]{} \
\dynkin[fold]{D}[1]{} \
```



We have to be careful about the 4-ply foldings of $D_{2\ell}^{(1)}$, for which we can have two different patterns, so by default, the package only draws as much as it can without distinguishing the two:

Default $D_{2\ell}^{(1)}$ and the two ways to finish it

```
\dynkin[ply=4]{D}[1]{****.*****.*****}%
\
\begin{dynkinDiagram}[ply=4]{D}[1]{****.*****.*****}%
\dynkinFold[bend right=90]{1}{13}%
\dynkinFold[bend right=90]{0}{14}%
\end{dynkinDiagram}%
\begin{dynkinDiagram}[ply=4]{D}[1]{****.*****.*****}%
\dynkinFold{0}{1}%
\dynkinFold{1}{13}%
\dynkinFold{13}{14}%
\end{dynkinDiagram}
```




Table 19: Some foldings of Dynkin diagrams. For these diagrams, we want to compare a folding diagram with the diagram that results when we fold it, so it looks best to set `fold radius` and `edge length` to equal lengths.

A_3	
C_2	

<code>\dynkin[fold]{A}[0]{3}</code>
<code>\dynkin{C}[0]{2}</code>









continued ...

Table 19: ...continued

$A_{2\ell-1}$		<code>\dynkin[fold]{A}{**.*****.**}</code>
C_ℓ		<code>\dynkin{C}{}</code>
B_3		<code>\dynkin[fold]{B}{0}{3}</code>
G_2		<code>\dynkin[reverse arrows]{G}{0}{2}</code>
D_4		<code>\dynkin[ply=3,fold right]{D}{4}</code>
G_2		<code>\dynkin{G}{2}</code>
$D_{\ell+1}$		<code>\dynkin[fold]{D}{}</code>
B_ℓ		<code>\dynkin{B}{}</code>
E_6		<code>\dynkin[fold]{E}{0}{6}</code>
F_4		<code>\dynkin[reverse arrows]{F}{0}{4}</code>
A_3^1		<code>\dynkin[ply=4]{A}{1}{3}</code>
A_1^1		<code>\dynkin{A}{1}{1}</code>
$A_{2\ell-1}^1$		<code>\dynkin[fold]{A}{1}{**.*****.**}</code>
C_ℓ^1		<code>\dynkin{C}{1}</code>
B_3^1		<code>\dynkin[ply=3]{B}{1}{3}</code>
A_2^2		<code>\dynkin{A}{2}{2}</code>
B_3^1		<code>\dynkin[ply=2]{B}{1}{3}</code>
G_2^1		<code>\dynkin{G}{1}{2}</code>
B_ℓ^1		<code>\dynkin[fold]{B}{1}</code>
D_ℓ^2		<code>\dynkin{D}{2}</code>
D_4^1		<code>\dynkin[ply=3]{D}{1}{4}</code>
B_3^1		<code>\dynkin{B}{1}{3}</code>

continued ...

Table 19: ... continued

D_4^1		<code>\dynkin[ply=3]{D}{1}{4}</code>
G_2^1		<code>\dynkin[G]{1}{2}</code>
$D_{\ell+1}^1$		<code>\dynkin[fold]{D}{1}{}</code>
D_{ℓ}^2		<code>\dynkin[D]{2}{}</code>
$D_{\ell+1}^1$		<code>\dynkin[fold right]{D}{1}{}</code>
B_{ℓ}^1		<code>\dynkin[B]{1}{}</code>
$D_{2\ell}^1$		<code>\begin{dynkinDiagram}[ply=4]{D}{1}% {****,*****,*} \dynkinFold{0}{1} \dynkinFold{1}{13} \dynkinFold{13}{14} \end{dynkinDiagram}</code>
A_{odd}^2		<code>\dynkin[A]{2}{odd}</code>
$D_{2\ell}^1$		<code>\begin{dynkinDiagram}[ply=4]{D}{1}% {****,*****,*} \dynkinFold[bend right=90]{1}{13} \dynkinFold[bend right=90]{0}{14} \end{dynkinDiagram}</code>
A_{even}^2		<code>\dynkin[A]{2}{even}</code>
E_6^1		<code>\dynkin[fold]{E}{1}{6}</code>
F_4^1		<code>\dynkin[reverse arrows]{F}{1}{4}</code>
E_6^1		<code>\dynkin[ply=3]{E}{1}{6}</code>
D_4^3		<code>\dynkin[D]{3}{4}</code>
E_7^1		<code>\dynkin[fold]{E}{1}{7}</code>
E_6^2		<code>\dynkin[E]{2}{6}</code>

continued ...

Table 19: ...continued

F_4^1		<code>\dynkin{F}{1}{4}</code>
G_2^1		<code>\dynkin{G}{1}{2}</code>
A_{odd}^2		<code>\dynkin[odd,fold]{A}{2}{****.***}</code>
A_{even}^2		<code>\dynkin{A}{2}{even}</code>
D_3^2		<code>\dynkin{D}{2}{3}</code>
A_2^2		<code>\dynkin{A}{2}{2}</code>


Table 20: Frobenius fixed point subgroups of finite simple groups of Lie type [4] p. 15

$A_{\ell \geq 1}$		<code>\dynkin{A}{}{}</code>
${}^2A_{\ell \geq 2}$		<code>\dynkin{fold}{A}{}{}</code>
$B_{\ell \geq 2}$		<code>\dynkin{B}{}{}</code>
2B_2		<code>\dynkin{fold}{B}{2}</code>
$C_{\ell \geq 3}$		<code>\dynkin{C}{}{}</code>
$D_{\ell \geq 4}$		<code>\dynkin{D}{}{}</code>
${}^2D_{\ell \geq 4}$		<code>\dynkin{fold}{D}{}{}</code>
3D_4		<code>\dynkin{ply=3}{D}{4}</code>
E_6		<code>\dynkin{E}{6}{}</code>
2E_6		<code>\dynkin{fold}{E}{6}</code>
E_7		<code>\dynkin{E}{7}{}</code>
E_8		<code>\dynkin{E}{8}{}</code>
F_4		<code>\dynkin{F}{4}{}</code>
2F_4		<code>\dynkin{fold}{F}{4}</code>
G_2		<code>\dynkin{G}{2}{}</code>
2G_2		<code>\dynkin{fold}{G}{2}</code>

24. ROOT ORDERING

Root ordering

```
\dynkin[label,ordering=Adams]{E}{6}
\dynkin[label,ordering=Bourbaki]{E}{6}
\dynkin[label,ordering=Carter]{E}{6}
\dynkin[label,ordering=Dynkin]{E}{6}
\dynkin[label,ordering=Kac]{E}{6}
```



Default is Bourbaki. Sources are Adams [1] p. 56–57, Bourbaki [3] p. pp. 265–290 plates I–IX, Carter [5] p. 540–609, Dynkin [8], Kac [15] p. 43.

	Adams	Bourbaki	Carter	Dynkin	Kac
E_6					
E_7					
E_8					
F_4					
G_2					

The marks are set down in order according to the current root ordering:

```
\dynkin[label]{E}{*otxXOt*}
\dynkin[label,ordering=Carter]{E}{*otxXOt*}
\dynkin[label,ordering=Kac]{E}{*otxXOt*}
```



25. CONNECTING DYNKIN DIAGRAMS

We can make some sophisticated folded diagrams by drawing multiple diagrams, each with a name:

Name a diagram

```
\dynkin[name=Bob]{D}{6}
```



We can then connect the two with folding edges:

Connect diagrams

```
\begin{dynkinDiagram}[name=upper]{A}{3}
    \node (current) at ($(upper root 1)+(0,-.3cm)$) {};
    \dynkin[at=(current),name=lower]{A}{3}
    \begin{scope}[on background layer]
        \foreach \i in {1,...,3}%
        {
            \draw[/Dynkin diagram/fold style]
                ($({upper root \i})$)
                -- ($({lower root \i})$);%
        }
    \end{scope}
\end{dynkinDiagram}
```




The following diagrams arise in the Satake diagrams of the pseudo-Riemannian symmetric spaces [2].


```
\pgfkeys{/Dynkin diagram,edge length=.5cm,fold radius=.5cm}
\begin{tikzpicture}
    \dynkin[name=1]{A}{IIIb}
    \node (a) at (-.3,-.4){};
    \dynkin[name=2,at=(a)]{A}{IIIb}
    \begin{scope}[on background layer]
        \foreach \i in {1,...,7}%
        {
            \draw[/Dynkin diagram/fold style]
                ($({1 root \i})$)
                --
                ($({2 root \i})$);%
        }
    \end{scope}
\end{tikzpicture}
```



```
\pgfkeys{/Dynkin diagram,
edge length=.75cm,
edge/.style={draw=example-color,double=black,very thick}}
\begin{tikzpicture}
\foreach \d in {1,...,4}
{
    \node (current) at ($(\d*.05,\d*.3)$){};
    \dynkin[name=\d,at=(current)]{D}{oo.oooo}
}
\begin{scope}[on background layer]
\foreach \i in {1,...,6}%
{%
    \draw[/Dynkin diagram/fold style] ($(1 root \i)$) -- ($(2
root \i)$);%
    \draw[/Dynkin diagram/fold style] ($(2 root \i)$) -- ($(3
root \i)$);%
    \draw[/Dynkin diagram/fold style] ($(3 root \i)$) -- ($(4
root \i)$);%
}
\end{scope}
\end{tikzpicture}
```




26. OTHER EXAMPLES




```
\tikzset{/Dynkin diagram,edge length=1cm,fold radius=1cm}
\tikzset{/Dynkin diagram,label macro/.code={\alpha_{\#1}},label macro*/.code={\beta_{\#1}}}
\({}^1 D_4\)\ 4-ply tied straight:
\begin{dynkinDiagram}[ply=4]{D}{1}%
{****.*****.****}
\dynkinFold{0}{1}
\dynkinFold{1}{13}
\dynkinFold{13}{14}
\dynkinLabelRoots{0,...,14}
\dynkinLabelRoots*{0,...,14}
\end{dynkinDiagram}
\({}^1 D_4\)\ 4-ply tied bending:
\begin{dynkinDiagram}[ply=4]{D}{1}%
{****.*****.****}
\dynkinFold{1}{13}
\dynkinFold[bend right=65]{0}{14}
\dynkinLabelRoots{0,...,14}
```

```
\dynkinLabelRoots*{0,...,14}
\end{dynkinDiagram}
```


Below we draw the Vogan diagrams of some affine Lie superalgebras [21, 20].




```
\dynkin[label,fold]{D}{1}{oo.oto.ooo}
```


 $\mathfrak{sl}(2m+1|2n)^2$


```
\dynkin[label]{B}{1}{oo.oto.oo}
```




```
\dynkin[label]{B}{1}{oO.oto.oO}
```




```
\dynkin[label,fold]{B}{1}{oo.oto.oo}
```








 $\mathfrak{sl}(2m+1|2n+1)^2$

```
\dynkin[label]{D}{2}{o.oto.oo}
```




```
\dynkin[label]{D}{2}{o.OtO.oo}
```




$\mathfrak{sl}(2|2n+1)^{(2)}$
 $\backslash \text{dynkin}[ply=2, \text{label}, \text{double edges}]\{\text{B}\}[1]\{\text{oo.Oto.Oo}\}$

 $\backslash \text{dynkin}[ply=2, \text{label}, \text{double fold}]\{\text{B}\}[1]\{\text{oo.Oto.Oo}\}$

 $\backslash \text{dynkin}[ply=2, \text{label}, \text{double edges}]\{\text{B}\}[1]\{\text{oo.OtO.oo}\}$

 $\backslash \text{dynkin}[ply=2, \text{label}, \text{double fold}]\{\text{B}\}[1]\{\text{oo.OtO.oo}\}$

 $\mathfrak{sl}(2|2n)^{(2)}$
 $\backslash \text{dynkin}[ply=2, \text{label}, \text{double edges}]\{\text{D}\}[1]\{\text{oo.oto.ooo}\}$

 $\backslash \text{dynkin}[ply=2, \text{label}, \text{double fold left}]\{\text{D}\}[1]\{\text{oo.oto.ooo}\}$


$\mathfrak{osp}(2m|2n)^{(2)}$


```
\dynkin[label,label macro/.code={\begin{array}{c} \text{\tiny \texttt{D}} \\[-1ex] \text{\tiny \texttt{o.oto.oo}} \end{array}}]
```




```
\dynkin[label,label macro/.code={\begin{array}{c} \text{\tiny \texttt{D}} \\[-1ex] \text{\tiny \texttt{o.Oto.Oo}} \end{array}}]
```


 $\mathfrak{osp}(2|2n)^{(2)}$


```
\dynkin[label,label macro/.code={\begin{array}{c} \text{\tiny \texttt{D}} \\[-1ex] \text{\tiny \texttt{affine mark=*\{o.o.o.o*}} \end{array}}]
```




```
\dynkin[label,label macro/.code={\begin{array}{c} \text{\tiny \texttt{D}} \\[-1ex] \text{\tiny \texttt{affine mark=*\{o.O.o.o*}} \end{array}}]
```


 $\mathfrak{sl}(1|2n+1)^4$

```
\dynkin[label,label macro/.code={\begin{array}{c} \text{\tiny \texttt{D}} \\[-1ex] \text{\tiny \texttt{o.o.o.o*}} \end{array}}]
```



```
\dynkin[label,label macro/.code={\{1\}}{\{D\}}{\{2\}}{\{o.o.O.o*\}}
```




A^1


```
\begin{tikzpicture}
\dynkin[name=upper]{A}{oo.t.oo}
\node (Dynkin current) at (upper root 1){};
\dynkinSouth
\dynkin[at=(Dynkin current),name=lower]{A}{oo.t.oo}
\begin{scope}[on background layer]
\foreach \i in {1,...,5}{
    \draw[/Dynkin diagram/fold style]
    ($(\text{upper root }\i)$) -- ($(\text{lower root }\i)$);
}
\end{scope}
\end{tikzpicture}
```




```
\dynkin[fold]{A}{1}{oo.t.ooooo.t.oo}
```



```
\dynkin[fold,affine mark=t]{A}{1}{oo.o.ootoo.o.oo}
```




```
\dynkin[affine mark=t]{A}{1}{o*.t.*o}
```




B^1


```
\dynkin[affine mark=*]{A}[2]{o.oto.o*}
```



```
\dynkin[affine mark=*]{A}[2]{o.oto.o*}
```




```
\dynkin[affine mark=*]{A}[2]{o.ooo.oo}
```




```
\dynkin[odd]{A}[2]{oo.*to.*o}
```



```
\dynkin[odd,fold]{A}[2]{oo.oto.oo}
```



```
\dynkin[odd,fold]{A}[2]{o*.oto.o*}
```

 D^1

```
\dynkin{D}{otoo}
```



```
\dynkin{D}{ot*o}
```




```
\dynkin[fold]{D}{otoo}
```




C^1

```
\dynkin[double edges,fold,affine mark=t,odd]{A}[2]{to.o*}
```



```
\dynkin[double edges,fold,affine mark=t,odd]{A}[2]{t*.oo}
```



F^1

```
\begin{dynkinDiagram}{A}{oto*}%
  \dynkinQuadrupleEdge{1}{2}%
  \dynkinTripleEdge{4}{3}%
\end{dynkinDiagram}%
```



```
\begin{dynkinDiagram}{A}{*too}%
  \dynkinQuadrupleEdge{1}{2}%
  \dynkinTripleEdge{4}{3}%
\end{dynkinDiagram}%
```



G^1

```
\begin{dynkinDiagram}{A}{ot*oo}%
\dynkinQuadrupleEdge{1}{2}%
\dynkinDefiniteDoubleEdge{4}{3}%
\end{dynkinDiagram}%
```

```
\begin{dynkinDiagram}{A}{oto*o}%
\dynkinQuadrupleEdge{1}{2}%
\dynkinDefiniteDoubleEdge{4}{3}%
\end{dynkinDiagram}%
```

```
\begin{dynkinDiagram}{A}{*too*}%
\dynkinQuadrupleEdge{1}{2}%
\dynkinDefiniteDoubleEdge{4}{3}%
\end{dynkinDiagram}%
```

```
\begin{dynkinDiagram}{A}{*tooo}%
\dynkinQuadrupleEdge{1}{2}%
\dynkinDefiniteDoubleEdge{4}{3}%
\end{dynkinDiagram}%
```

27. EXAMPLE: THE COMPLEX SIMPLE LIE ALGEBRAS

\mathfrak{g}	Diagram	Weights	Roots	Simple roots
A_n		$\frac{1}{r+1}\mathbb{Z}^{n+1} / \langle \sum e_j \rangle$	$e_i - e_j$	$e_i - e_{i+1}$
B_n		$\frac{1}{2}\mathbb{Z}^n$	$\pm e_i, \pm e_i \pm e_j, i \neq j$	$e_i - e_{i+1}, e_n$
C_n		\mathbb{Z}^n	$\pm 2e_i, \pm e_i \pm e_j, i \neq j$	$e_i - e_{i+1}, 2e_n$
D_n		$\frac{1}{2}\mathbb{Z}^n$	$\pm e_i \pm e_j, i \neq j$	$e_i - e_{i+1}, \quad i \leq n-1$ $e_{n-1} + e_n$

\mathfrak{g}	Diagram	Weights	Roots	Simple roots
E_8		$\frac{1}{2}\mathbb{Z}^8$	$\pm 2e_i \pm 2e_j, \quad i \neq j,$ $\sum_i (-1)^{m_i} e_i, \quad \sum m_i \text{ even}$	$2e_1 - 2e_2,$ $2e_2 - 2e_3,$ $2e_3 - 2e_4,$ $2e_4 - 2e_5,$ $2e_5 - 2e_6,$ $2e_6 + 2e_7,$ $- \sum e_j,$ $2e_6 - 2e_7$
E_7		$\frac{1}{2}\mathbb{Z}^8 / \langle e_1 - e_2 \rangle$	quotient of E_8	quotient of E_8
E_6		$\frac{1}{3}\mathbb{Z}^8 / \langle e_1 - e_2, e_2 - e_3 \rangle$	quotient of E_8	quotient of E_8
F_4		\mathbb{Z}^4	$\pm 2e_i,$ $\pm 2e_i \pm 2e_j, \quad i \neq j,$ $\pm e_1 \pm e_2 \pm e_3 \pm e_4$	$2e_2 - 2e_3,$ $2e_3 - 2e_4,$ $2e_4,$ $e_1 - e_2 - e_3 - e_4$
G_2		$\mathbb{Z}^3 / \langle \sum e_j \rangle$	$\pm(1, -1, 0),$ $\pm(-1, 0, 1),$ $\pm(0, -1, 1),$ $\pm(2, -1, -1),$ $\pm(1, -2, 1),$ $\pm(-1, -1, 2)$	$(-1, 0, 1),$ $(2, -1, -1)$


```

\NewDocumentEnvironment{bunch}{}%
{\renewcommand*{\arraystretch}{1}\begin{array}{@{}l|l@{}}\hline & \\ \midrule}{\hline \midrule\end{array}}
\small
\NewDocumentCommand{\nct}[2]{\newcolumntype{#1}{>{\columncolor[gray]{.9}>{$\cdot$}m{#2cm}<{$\cdot$}}}}{\nct{G}{.3}\nct{D}{2.1}\nct{W}{3}\nct{R}{3.7}\nct{S}{3}}
\NewDocumentCommand{\LieG}{\mathfrak{g}}{\mathfrak{g}}
\NewDocumentCommand{\Wom}{\mathbf{Z}}{\mathbf{Z}^{\#2}\IfValueT{\#1}{/\left<\#1\right>}}
\renewcommand*{\arraystretch}{1.5}
\NewDocumentCommand{\quo}{\text{quotient of }}{E_8}
\begin{longtable}{@{}GDWRS@{}}
\LieG&\text{Diagram}&\text{Weights}&\text{Roots}&\text{Simple roots}\\ \hline
\endfirsthead
\LieG&\text{Diagram}&\text{Weights}&\text{Roots}&\text{Simple roots}\\ \hline
A_n &\dynkin{A}{}&&&\\
B_n &\dynkin{B}{}&&&\\
C_n &\dynkin{C}{}&&&\\
D_n &\dynkin{D}{}&&&\\
E_8 &\dynkin{E}{8}&&&\\
\begin{bunch}e_{i+1},&i\leq n-1\end{bunch}&&&\\
2e_1-2e_2,\&2e_2-2e_3,\&2e_3-2e_4,\&2e_4-2e_5,\&2e_5-2e_6,\&2e_6+2e_7,\&\\
-\sum e_j,\&2e_6-2e_7\\ \end{bunch}\\ \end{longtable}

```

```
E_7&\dynkin{E}{7}&\frac{1}{2}\W[e_1-e_2]{8}&\quo&\quo\\
E_6&\dynkin{E}{6}&\frac{1}{3}\W[e_1-e_2,e_2-e_3]{8}&\quo&\quo\\
F_4&\dynkin{F}{4}&\W{4}&
\begin{bunch}\pm 2e_i,\pm 2e_i \pm 2e_j, \quad i \neq j, \pm e_1 \pm e_2 \pm e_3 \pm e_4
\end{bunch}&
\begin{bunch}2e_2-2e_3, 2e_3-2e_4, 2e_4, e_1-e_2-e_3-e_4
\end{bunch}\\
G_2&\dynkin{G}{2}&\W[\sum e_j]{3}&
\begin{bunch}
\pm(1,-1,0), \pm(-1,0,1), \pm(0,-1,1), \pm(2,-1,-1), \pm(1,-2,1), \pm(-1,-1,2)
\end{bunch}&
\begin{bunch}(-1,0,1), (2,-1,-1)
\end{bunch}
\end{longtable}
```

28. AN EXAMPLE OF MIKHAIL BOROVOI



```
\tikzset{big arrow/.style={-Stealth, line cap=round, line width=1mm,
shorten <=1mm, shorten >=1mm}}
\newcommand{\catholic}[2]{\draw[big arrow,green!25!white]
(root #1) to (root #2);}
\newcommand{\protestant}[2]{
\begin{scope}[transparency group, opacity=.25]
\draw[big arrow,orange] (root #1) to (root #2);
\end{scope}}
\begin{dynkinDiagram}[edge length=1.2cm,
indefinite edge/.style={thick,loosely dotted},
labels*={0,1,2,3,\ell-3,\ell-2,\ell-1,\ell}]{D}{1}{}
\catholic{0}{6}\catholic{1}{7}
\protestant{7}{0}\protestant{6}{1}
\end{dynkinDiagram}
```

29. SYNTAX

The syntax is `\dynkin[<options>]{<letter>}[<twisted rank>]{<rank>}` where `<letter>` is A, B, C, D, E, F or G, the family of root system for the Dynkin diagram, `<twisted rank>` is 0, 1, 2, 3 (default is 0) representing:

- 0 finite root system
- 1 affine extended root system, i.e. of type ⁽¹⁾
- 2 affine twisted root system of type ⁽²⁾
- 3 affine twisted root system of type ⁽³⁾

and `<rank>` is

- (1) an integer representing the rank or
- (2) blank to represent an indefinite rank or
- (3) the name of a Satake diagram as in section 4.

The environment syntax is `\begin{dynkinDiagram}` followed by the same parameters as `\dynkin`, then various Dynkin diagram and TikZ commands, and then `\end{dynkinDiagram}`.

30. OPTIONS

```
ceref = <true or false>,
default : false
    whether to draw roots in a “ceref” style.
edge length = <number>cm,
default : .35cm
    distance between nodes in the Dynkin diagram
edge/.style = TikZ style data,
default : solid,draw=black,fill=white,thin
    style of edges in the Dynkin diagram
Kac = <true or false>,
default : false
    whether to draw in the style of [15]
name = <string>,
default : anonymous
    A name for the Dynkin diagram, with anonymous treated as a
    blank; see section 25.
parabolic = <integer>,
default : 0
    A parabolic subgroup with specified integer, where the integer
    is computed as  $n = \sum 2^{i-1}a_i$ ,  $a_i = 0$  or 1, to say that root  $i$  is
    crossed, i.e. a noncompact root.
root radius = <number>cm,
default : .05cm
    size of the dots and of the crosses in the Dynkin diagram
text style = <TikZ style data>,
default : scale=.7
    Style for any labels on the roots.
mark = <o,O,t,x,X,*>,
    continued ...
```

Table 23: ... continued

```

default : *
    default root mark
affine mark = o,O,t,x,X,*,
default : *
    default root mark for root zero in an affine Dynkin diagram
label = true or false,
default : false
    whether to label the roots according to the current labelling scheme.
label macro = <1-parameter TEX macro>,
default : #1
    the current labelling scheme for roots.
label macro* = <1-parameter TEX macro>,
default : #1
    the current labelling scheme for alternate roots.
label height = <1-parameter TEX macro>,
default : b
    the current maximal height of text labels for the roots, set by
    giving mathematics text of that height.
label depth = <1-parameter TEX macro>,
default : g
    the current maximal depth of text labels for the roots, set by
    giving mathematics text of that depth.
make indefinite edge = <edge pair i-j or list of such>,
default : {}
    edge pair or list of edge pairs to treat as having indefinitely many
    roots on them.
indefinite edge ratio = <float>,
default : 1.6
    ratio of indefinite edge lengths to other edge lengths.
indefinite edge/.style = <TikZ style data>,
default : solid,draw=black,fill=white,thin,densely dotted
    style of the dotted or dashed middle third of each indefinite edge.
backwards = <true or false>,
default : false
    whether to reverse right to left.
upside down = <true or false>,
default : false
    whether to reverse up to down.
arrows = <true or false>,
default : true
    whether to draw the arrows that arise along the edges.
reverse arrows = <true or false>,
default : true
    whether to reverse the direction of the arrows that arise along the
    edges.
fold = <true or false>,

```

continued ...

Table 23: ... continued

```

default : true
    whether, when drawing Dynkin diagrams, to draw them 2-ply.
ply = ⟨0,1,2,3,4⟩,
default : 0
    how many roots get folded together, at most.
fold left = ⟨true or false⟩,
default : true
    whether to fold the roots on the left side of a Dynkin diagram.
fold right = ⟨true or false⟩,
default : true
    whether to fold the roots on the right side of a Dynkin diagram.
fold radius = ⟨length⟩,
default : .3cm
    the radius of circular arcs used in curved edges of folded Dynkin
    diagrams.
fold style/.style = ⟨TikZ style data⟩,
default : solid,draw=black!40,fill=none,line width=radius
    when drawing folded diagrams, style for the fold indicators.
*/.style = ⟨TikZ style data⟩,
default : solid,draw=black,fill=black
    style for roots like •
o/.style = ⟨TikZ style data⟩,
default : solid,draw=black,fill=black
    style for roots like o
O/.style = ⟨TikZ style data⟩,
default : solid,draw=black,fill=black
    style for roots like @
t/.style = ⟨TikZ style data⟩,
default : solid,draw=black,fill=black
    style for roots like °
x/.style = ⟨TikZ style data⟩,
default : solid,draw=black,line cap=round
    style for roots like x
X/.style = ⟨TikZ style data⟩,
default : solid,draw=black,thick,line cap=round
    style for roots like x
fold left style/.style = ⟨TikZ style data⟩,
default :
    style to override the fold style when folding roots together on the
    left half of a Dynkin diagram
fold right style/.style = ⟨TikZ style data⟩,
default :
    style to override the fold style when folding roots together on the
    right half of a Dynkin diagram
double edges = ⟨⟩,
default : not set

```

continued ...

Table 23: ...continued

set to override the <code>fold</code> style when folding roots together in a Dynkin diagram, so that the foldings are indicated with double edges (like those of an F_4 Dynkin diagram without arrows).
<code>double fold = </>,</code> default : not set
set to override the <code>fold</code> style when folding roots together in a Dynkin diagram, so that the foldings are indicated with double edges (like those of an F_4 Dynkin diagram without arrows), but filled in solidly.
<code>double left = </>,</code> default : not set
set to override the <code>fold</code> style when folding roots together at the left side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an F_4 Dynkin diagram without arrows).
<code>double fold left = </>,</code> default : not set
set to override the <code>fold</code> style when folding roots together at the left side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an F_4 Dynkin diagram without arrows), but filled in solidly.
<code>double right = </>,</code> default : not set
set to override the <code>fold</code> style when folding roots together at the right side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an F_4 Dynkin diagram without arrows).
<code>double fold right = </>,</code> default : not set
set to override the <code>fold</code> style when folding roots together at the right side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an F_4 Dynkin diagram without arrows), but filled in solidly.
<code>arrow color = </>,</code> default : black
set to override the default color for the arrows in nonsimply laced Dynkin diagrams.
<code>Coxeter = <true or false>,</code> default : false
whether to draw a Coxeter diagram, rather than a Dynkin diagram.
<code>ordering = <Adams, Bourbaki, Carter, Dynkin, Kac>,</code> default : Bourbaki
which ordering of the roots to use in exceptional root systems as in section 24.

All other options are passed to TikZ.

REFERENCES

1. J. F. Adams, *Lectures on exceptional Lie groups*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1996, With a foreword by J. Peter May, Edited by Zafer Mahmud and Mamoru Mimura. MR 1428422
2. Kurando Baba, *Satake diagrams and restricted root systems of semisimple pseudo-Riemannian symmetric spaces*, Tokyo J. Math. **32** (2009), no. 1, 127–158. MR 2541161
3. Nicolas Bourbaki, *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressley. MR 1890629
4. R. W. Carter, *On the representation theory of the finite groups of Lie type over an algebraically closed field of characteristic o* [MR1170353 (93j:20034)], Algebra, IX, Encyclopaedia Math. Sci., vol. 77, Springer, Berlin, 1995, pp. 1–120, 235–239. MR 1392478
5. ———, *Lie algebras of finite and affine type*, Cambridge Studies in Advanced Mathematics, vol. 96, Cambridge University Press, Cambridge, 2005. MR 2188930
6. Meng-Kiat Chuah, *Cartan automorphisms and Vogan superdiagrams*, Math. Z. **273** (2013), no. 3–4, 793–800. MR 3030677
7. Cristina Draper Fontanals and Valerio Guido, *On the real forms of the exceptional Lie algebra \mathfrak{e}_6 and their Satake diagrams*, Non-associative and non-commutative algebra and operator theory, Springer Proc. Math. Stat., vol. 160, Springer, Cham, 2016, pp. 211–226. MR 3613831
8. E. B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, Mat. Sbornik N.S. **30(72)** (1952), 349–462 (3 plates), Reprinted in English translation in [9]. MR 0047629
9. ———, *Selected papers of E. B. Dynkin with commentary*, American Mathematical Society, Providence, RI; International Press, Cambridge, MA, 2000, Edited by A. A. Yushkevich, G. M. Seitz and A. L. Onishchik. MR 1757976
10. L. Frappat, A. Sciarrino, and P. Sorba, *Structure of basic Lie superalgebras and of their affine extensions*, Comm. Math. Phys. **121** (1989), no. 3, 457–500. MR 990776
11. William Fulton and Joe Harris, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991, A first course, Readings in Mathematics. MR 1153249
12. L. C. Grove and C. T. Benson, *Finite reflection groups*, second ed., Graduate Texts in Mathematics, vol. 99, Springer-Verlag, New York, 1985. MR 777684
13. Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics, vol. 34, American Mathematical Society, Providence, RI, 2001, Corrected reprint of the 1978 original. MR 1834454
14. James E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, vol. 29, Cambridge University Press, Cambridge, 1990. MR 1066460
15. Victor G. Kac, *Infinite-dimensional Lie algebras*, third ed., Cambridge University Press, Cambridge, 1990. MR 1104219
16. S. Pratik Khastgir and Ryu Sasaki, *Non-canonical folding of Dynkin diagrams and reduction of affine Toda theories*, Progr. Theoret. Phys. **95** (1996), no. 3, 503–518. MR 1388245
17. Robert P. Langlands, *Euler products*, Yale University Press, New Haven, Conn.-London, 1971, A James K. Whittemore Lecture in Mathematics given at Yale University, 1967, Yale Mathematical Monographs, 1. MR 0419366
18. A. L. Onishchik and È. B. Vinberg, *Lie groups and algebraic groups*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1990, Translated from the Russian and with a preface by D. A. Leites. MR 91g:22001
19. A. L. Onishchik and È. B. Vinberg, *Lie groups and algebraic groups*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1990, Translated from the Russian and with a preface by D. A. Leites. MR 1064110
20. B. Ransingh, *Vogan diagrams of affine twisted Lie superalgebras*, ArXiv e-prints (2013), 1–9.
21. Biswajit Ransingh, *Vogan diagrams of untwisted affine Kac-Moody superalgebras*, Asian-Eur. J. Math. **6** (2013), no. 4, 1350062, 10. MR 3149279
22. V. Regelskis and B. Vlaar, *Reflection matrices, coideal subalgebras and generalized Satake diagrams of affine type*, ArXiv e-prints (2016), 1–118.
23. Ichirô Satake, *Algebraic structures of symmetric domains*, Kanô Memorial Lectures, vol. 4, Iwanami Shoten, Tokyo; Princeton University Press, Princeton, N.J., 1980. MR 591460
24. T. A. Springer, *Linear algebraic groups*, second ed., Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2009. MR 2458469

25. È. B. Vinberg (ed.), *Lie groups and Lie algebras, III*, Encyclopaedia of Mathematical Sciences, vol. 41, Springer-Verlag, Berlin, 1994, Structure of Lie groups and Lie algebras, A translation of it Current problems in mathematics. Fundamental directions. Vol. 41 (Russian), Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1990 [MR1056485 (91b:22001)], Translation by V. Minachin [V. V. Minakhin], Translation edited by A. L. Onishchik and È. B. Vinberg. MR 1349140
26. Jean-Bernard Zuber, *Generalized Dynkin diagrams and root systems and their folding*, Topological field theory, primitive forms and related topics (Kyoto, 1996), Progr. Math., vol. 160, Birkhäuser Boston, Boston, MA, 1998, pp. 453–493. MR 1653035

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE CORK, CORK, IRELAND
Email address: b.mckay@ucc.ie