

# THE DYNKIN DIAGRAMS PACKAGE

## VERSION 3.12

BEN MCKAY

### CONTENTS

1. Quick introduction, 1
2. Set options globally, 3
3. Coxeter diagrams, 3
4. Satake diagrams, 4
5. Labels for the roots, 5
6. Style, 7
7. Suppress or reverse arrows, 8
8. Drawing on top of a Dynkin diagram, 9
9. Mark lists, 9
10. Indefinite edges, 10
11. Parabolic subgroups, 13
12. Extended Dynkin diagrams, 15
13. Affine twisted and untwisted Dynkin diagrams, 15
14. Extended Coxeter diagrams, 17
15. Kac style, 18
16. Folded Dynkin diagrams, 19
17. Root ordering, 24
18. Connecting Dynkin diagrams, 24
19. Other examples, 26
20. Syntax, 35
21. Options, 35
- References, 38

### 1. QUICK INTRODUCTION

Load the Dynkin diagram package (see options below)

```
\documentclass{amsart}
\usepackage{dynkin-diagrams}
\begin{document}
The Dynkin diagram of \(\mathbf{B}_3\) is \dynkin{B}{3}.
\end{document}
```

Invoke it

```
The Dynkin diagram of \(\mathbf{B}_3\) is \dynkin{B}{3}.
```

The Dynkin diagram of  $B_3$  is  $\bullet\rightarrowtail\bullet$ .

Inside a *TikZ* statement

```
The Dynkin diagram of \(\mathbf{B}_3\) is  
\tikz[baseline=-0.5ex] \dynkin{B}{3};
```

The Dynkin diagram of  $B_3$  is  $\bullet\rightarrowtail\bullet$

Inside a *TikZ* environment

```
The Dynkin diagram of \(\mathbf{B}_3\) is  
\begin{tikzpicture}[baseline=-0.5ex]  
  \dynkin{B}{3}  
\end{tikzpicture}
```

The Dynkin diagram of  $B_3$  is  $\bullet\rightarrowtail\bullet$

Indefinite rank Dynkin diagrams

```
\dynkin{B}{}
```

Table 1: The Dynkin diagrams of the reduced simple root systems [2] pp. 265–290, plates I–IX

$A_n$		<code>\dynkin{A}{}</code>
$C_n$		<code>\dynkin{C}{}</code>
$D_n$		<code>\dynkin{D}{}</code>
$E_6$		<code>\dynkin{E}{6}</code>
$E_7$		<code>\dynkin{E}{7}</code>
$E_8$		<code>\dynkin{E}{8}</code>
$F_4$		<code>\dynkin{F}{4}</code>
$G_2$		<code>\dynkin{G}{2}</code>

## 2. SET OPTIONS GLOBALLY

Most options set globally ...

```
\pgfkeys{/Dynkin diagram,edgeLength=.5cm,foldradius=.5cm}
```


... or pass to the package

```
\usepackage[
    ordering=Kac,
    edge/.style=blue,
    mark=o,
    radius=.06cm]
{dynkin-diagrams}
```

## 3. COXETER DIAGRAMS

Coxeter diagram option

```
\dynkin[Coxeter]{F}{4}
```



gonality option for  $G_2$  and  $I_n$  Coxeter diagrams

```
\(G_2=\dynkin[Coxeter,gonality=n]{G}{2}), \
\I_n=\dynkin[Coxeter,gonality=n]{I}{}\\
```

$G_2 = \bullet^n \bullet$ ,  $I_n = \bullet^n \bullet$

Table 2: The Coxeter diagrams of the simple reflection groups

$A_n$		<code>\dynkin[Coxeter]{A}{}</code>
$B_n$		<code>\dynkin[Coxeter]{B}{}</code>
$C_n$		<code>\dynkin[Coxeter]{C}{}</code>
$E_6$		<code>\dynkin[Coxeter]{E}{6}</code>
$E_7$		<code>\dynkin[Coxeter]{E}{7}</code>
$E_8$		<code>\dynkin[Coxeter]{E}{8}</code>
$F_4$		<code>\dynkin[Coxeter]{F}{4}</code>

continued ...


Table 2: ... continued

$G_2$		<code>\dynkin[Coxeter,gonality=n]{G}{2}</code>
$H_3$		<code>\dynkin[Coxeter]{H}{3}</code>
$H_4$		<code>\dynkin[Coxeter]{H}{4}</code>
$I_n$		<code>\dynkin[Coxeter,gonality=n]{I}{}{}</code>

## 4. SATAKE DIAGRAMS

Satake diagrams use the standard name instead of a rank

```
\\"(A_{IIIb})=\dynkin{A}{IIIb}\")
```



We use a solid gray bar to denote the folding of a Dynkin diagram, rather than the usual double arrow, since the diagrams turn out simpler and easier to read.

Table 3: The Satake diagrams of the real simple Lie algebras [10] p. 532–534

$A_I$		<code>\dynkin{A}{I}</code>
$A_{II}$		<code>\dynkin{A}{II}</code>
$A_{IIIa}$		<code>\dynkin{A}{IIIa}</code>
$A_{IIIb}$		<code>\dynkin{A}{IIIb}</code>
$A_{IV}$		<code>\dynkin{A}{IV}</code>
$B_I$		<code>\dynkin{B}{I}</code>
$B_{II}$		<code>\dynkin{B}{II}</code>
$C_I$		<code>\dynkin{C}{I}</code>
$C_{IIa}$		<code>\dynkin{C}{IIa}</code>
$C_{IIb}$		<code>\dynkin{C}{IIb}</code>
$D_{Ia}$		<code>\dynkin{D}{Ia}</code>
$D_{Ib}$		<code>\dynkin{D}{Ib}</code>

continued ...

Table 3: ... continued


$D_{Ic}$		<code>\dynkin{D}{Ic}</code>
$D_{II}$		<code>\dynkin{D}{II}</code>
$D_{IIIa}$		<code>\dynkin{D}{IIIa}</code>
$D_{IIIb}$		<code>\dynkin{D}{IIIb}</code>
$E_I$		<code>\dynkin{E}{I}</code>
$E_{II}$		<code>\dynkin{E}{II}</code>
$E_{III}$		<code>\dynkin{E}{III}</code>
$E_{IV}$		<code>\dynkin{E}{IV}</code>
$E_V$		<code>\dynkin{E}{V}</code>
$E_{VI}$		<code>\dynkin{E}{VI}</code>
$E_{VII}$		<code>\dynkin{E}{VII}</code>
$E_{VIII}$		<code>\dynkin{E}{VIII}</code>
$E_{IX}$		<code>\dynkin{E}{IX}</code>
$F_I$		<code>\dynkin{F}{I}</code>
$F_{II}$		<code>\dynkin{F}{II}</code>
$G_I$		<code>\dynkin{G}{I}</code>

## 5. LABELS FOR THE ROOTS

Label the roots by root number

`\dynkin[label]{B}{3}`

Make a macro to assign labels to roots

`\dynkin[label,labelMacro/.code={\alpha_{#1}}]{D}{5}`

## Label a single root

```
\begin{tikzpicture}
    \dynkin[B]{3}
    \dynkinLabelRoot{2}{\alpha_2}
\end{tikzpicture}
```



## Use a text style

```
\begin{tikzpicture}
    \dynkin[text/.style={scale=1.2}]{B}{3};
    \dynkinLabelRoot{2}{\alpha_2}
\end{tikzpicture}
```




## Access root labels via TikZ

```
\begin{tikzpicture}
    \dynkin[B]{3};
    \node[below] at (root 2) {\(\alpha_2)};
\end{tikzpicture}
```



## The labels have default locations

```
\begin{tikzpicture}
    \dynkin[E]{8};
    \dynkinLabelRoot{1}{\alpha_1}
    \dynkinLabelRoot{2}{\alpha_2}
    \dynkinLabelRoot{3}{\alpha_3}
\end{tikzpicture}
```



The starred form flips labels to alternate locations

```
\begin{tikzpicture}
    \dynkin[E]{8};
    \dynkinLabelRoot*[1]{\alpha_1}
    \dynkinLabelRoot*[2]{\alpha_2}
    \dynkinLabelRoot*[3]{\alpha_3}
\end{tikzpicture}
```



## 6. STYLE

Colours

```
\dynkin[edge/.style={blue!50,thick},*/.style=blue!50!red]{F}{4}
```




Edge lengths

```
\dynkin[edgeLength=1.2,parabolic=3]{A}{3}
```



Root marks

```
\dynkin[E]{8}
\dynkin[mark=*]{E}{8}
\dynkin[mark=o]{E}{8}
\dynkin[mark=0]{E}{8}
\dynkin[mark=t]{E}{8}
\dynkin[mark=x]{E}{8}
\dynkin[mark=X]{E}{8}
```




At the moment, you can only use:

- \* solid dot
- o hollow circle
- O double hollow circle
- t tensor root
- x crossed root
- X thickly crossed root

#### Mark styles

```
\dynkin[parabolic=124,x/.style={brown,very thick}]{E}{8}
```



#### Sizes of root marks

```
\dynkin[radius=.08cm,parabolic=3]{A}{3}
```



## 7. SUPPRESS OR REVERSE ARROWS


#### Some diagrams have double or triple edges

```
\dynkin{F}{4}
\dynkin{G}{2}
```



#### Suppress arrows

```
\dynkin[arrows=false]{F}{4}
\dynkin[arrows=false]{G}{2}
```



#### Reverse arrows

```
\dynkin[reverseArrows]{F}{4}
\dynkin[reverseArrows]{G}{2}
```



## 8. DRAWING ON TOP OF A DYNKIN DIAGRAM


TikZ can access the roots themselves

```
\begin{tikzpicture}
  \dynkin[A]{4};
  \fill[white,draw=black] (root 2) circle (.15cm);
  \fill[white,draw=black] (root 2) circle (.1cm);
  \draw[black] (root 2) circle (.05cm);
\end{tikzpicture}
```




Draw curves between the roots

```
\begin{tikzpicture}
  \dynkin[label]{E}{8}
  \draw[very thick, black!50,-latex]
    (root 3.south) to [out=-45, in=-135] (root 6.south);
\end{tikzpicture}
```



Change marks

```
\begin{tikzpicture}
  \dynkin[mark=o,label]{E}{8};
  \dynkinRootMark{*}{5}
  \dynkinRootMark{*}{8}
\end{tikzpicture}
```



## 9. MARK LISTS

The package allows a list of root marks instead of a rank:

A mark list

```
\dynkin{E}{oo**ttxx}
```



The mark list `oo**ttxx` has one mark for each root: `o`, `o`,  $\dots$ , `x`. Roots are listed in the current default ordering. (Careful: in an affine root system, a mark list will *not* contain a mark for root zero.)

Table 4: Classical Lie superalgebras [8]. We need a slightly larger radius parameter to distinguish the tensor product symbols from the solid dots.

$A_{mn}$		<code>\dynkin{A}{ooo.oto.oo}</code>
$B_{mn}$		<code>\dynkin{B}{ooo.oto.oo}</code>
$B_{0n}$		<code>\dynkin{B}{ooo.ooo.o*}</code>
$C_n$		<code>\dynkin{C}{too.oto.oo}</code>
$D_{mn}$		<code>\dynkin{D}{ooo.oto.oooo}</code>
$D_{21\alpha}$		<code>\dynkin{A}{oto}</code>
$F_4$		<code>\dynkin{F}{oot}</code>
$G_3$		<code>\dynkin[extended,affineMark=t]{G}{2}</code>

Table 5: Classical Lie superalgebras [8]. Here we see the problem with using the default radius parameter, which is too small for tensor product symbols.

$A_{mn}$		<code>\dynkin{A}{ooo.oto.oo}</code>
$B_{mn}$		<code>\dynkin{B}{ooo.oto.oo}</code>
$B_{0n}$		<code>\dynkin{B}{ooo.ooo.o*}</code>
$C_n$		<code>\dynkin{C}{too.oto.oo}</code>
$D_{mn}$		<code>\dynkin{D}{ooo.oto.oooo}</code>
$D_{21\alpha}$		<code>\dynkin{A}{oto}</code>
$F_4$		<code>\dynkin{F}{oot}</code>
$G_3$		<code>\dynkin[extended,affineMark=t]{G}{2}</code>

## 10. INDEFINITE EDGES

An *indefinite edge* is a dashed edge between two roots, indicating that an indefinite number of roots have been omitted from the Dynkin diagram. In between any two entries in a mark list, place a period to indicate an indefinite edge:

## Indefinite edges


```
\dynkin{D}{o.o.*.*.t.to.t}
```



In certain diagrams, roots may have an edge between them even though they are not subsequent in the ordering. For such rare situations, there is an option:


## Indefinite edge option

```
\dynkin[makeIndefiniteEdge={3-5},label]{D}{5}
```




## Give a list of edges to become indefinite

```
\dynkin[makeIndefiniteEdge/.list={1-2,3-5},label]{D}{5}
```



## Indefinite edge style

```
\dynkin[indefiniteEdge/.style={draw=black,fill=white,thin,densely
dashed},%
edgeLength=1cm,%
makeIndefiniteEdge={3-5}]
{D}{5}
```



The ratio of the lengths of indefinite edges to those of other edges

```
\dynkin[edgeLength = .5cm,%
       indefiniteEdgeRatio=3,%
       makeIndefiniteEdge={3-5}]
{D}{5}
```




Table 6: Springer's table of indices [20], pp. 320-321, with one form of  $E_7$  corrected

$A_n$		$d$	$n-d$
$A_n$		$d$	$rd$
$B_n$		$r$	
$C_n$		$d$	$rd$
$D_n$		$d$	$rd$
$E_6$			$\backslash\text{dynkin}\{\text{E}\}\{\astoooo\ast\}$
$E_6$			$\backslash\text{dynkin}\{\text{E}\}\{\circ*\circ*\circ o\}$
$E_6$			$\backslash\text{dynkin}\{\text{E}\}\{\circ*\circoooo\}$
$E_6$			$\backslash\text{dynkin}\{\text{E}\}\{**ooo*\}$
$E_7$			$\backslash\text{dynkin}\{\text{E}\}\{\astoooooo\}$
$E_7$			$\backslash\text{dynkin}\{\text{E}\}\{oooooo*\circ\}$
$E_7$			$\backslash\text{dynkin}\{\text{E}\}\{oooooo*\}$
$E_7$			$\backslash\text{dynkin}\{\text{E}\}\{\circoooo*\circ\}$
$E_7$			$\backslash\text{dynkin}\{\text{E}\}\{\astoooo**\}$
$E_7$			$\backslash\text{dynkin}\{\text{E}\}\{\ast o***o*\circ\}$
$E_8$			$\backslash\text{dynkin}\{\text{E}\}\{\astooooooo\}$

continued ...

Table 6: ... continued

$E_8$		<code>\dynkin{E}{ooooooo*}</code>
$E_8$		<code>\dynkin{E}{*oooooo*}</code>
$E_8$		<code>\dynkin{E}{oooooo**}</code>
$E_8$		<code>\dynkin{E}{*oooo***}</code>
$F_4$		<code>\dynkin{F}{ooo*}</code>
$D_4$		<code>\dynkin{D}{o*oo}</code>

## 11. PARABOLIC SUBGROUPS

Each set of roots is assigned a number, with each binary digit zero or one to say whether the corresponding root is crossed or not:

The flag variety of pointed lines in projective 3-space is associated to the Dynkin diagram `\dynkin[parabolic=3]{A}{3}`.




The flag variety of pointed lines in projective 3-space is associated to the Dynkin diagram  $\times \leftarrow \rightarrow \bullet$ .

Table 7: The Hermitian symmetric spaces

$A_n$		<code>\dynkin{A}{**.*x*.**}</code>
		Grassmannian of $k$ -planes in $\mathbb{C}^{n+1}$
$B_n$		<code>\dynkin[parabolic=1]{B}{}{}</code>
		$(2n-1)$ -dimensional hyperquadric, i.e. the variety of null lines in $\mathbb{C}^{2n+1}$
$C_n$		<code>\dynkin[parabolic=16]{C}{}{}</code>
		space of Lagrangian $n$ -planes in $\mathbb{C}^{2n}$

Table 7: continued ...

Table 7: ... continued

$D_n$		$\backslash\text{dynkin}[\text{parabolic}=1]\{\text{D}\}\{\}$
		( $2n - 2$ )-dimensional hyperquadric, i.e. the variety of null lines in $\mathbb{C}^{2n}$
$D_n$		$\backslash\text{dynkin}[\text{parabolic}=32]\{\text{D}\}\{\}$
		one component of the variety of maximal dimension null subspaces of $\mathbb{C}^{2n}$
$D_n$		$\backslash\text{dynkin}[\text{parabolic}=16]\{\text{D}\}\{\}$
		the other component
$E_6$		$\backslash\text{dynkin}[\text{parabolic}=1]\{\text{E}\}\{6\}$
		complexified octave projective plane
$E_6$		$\backslash\text{dynkin}[\text{parabolic}=32]\{\text{E}\}\{6\}$
		its dual plane
$E_7$		$\backslash\text{dynkin}[\text{parabolic}=64]\{\text{E}\}\{7\}$
		the space of null octave 3-planes in oc- tave 6-space

## 12. EXTENDED DYNKIN DIAGRAMS

Extended Dynkin diagrams

\dynkin[extended]{A}{7}



The extended Dynkin diagrams are also described in the notation of Kac [12] p. 55 as affine untwisted Dynkin diagrams: we extend `\dynkin{A}{7}` to become `\dynkin{A}{1}{7}`:

Extended Dynkin diagrams

\dynkin{A}{1}{7}




Table 8: The Dynkin diagrams of the extended simple root systems

$A_1^1$	$\circ\leftrightarrow\bullet$	<code>\dynkin[extended]{A}{1}</code>
$A_n^1$		<code>\dynkin[extended]{A}{()}</code>
$B_n^1$		<code>\dynkin[extended]{B}{()}</code>
$C_n^1$		<code>\dynkin[extended]{C}{()}</code>
$D_n^1$		<code>\dynkin[extended]{D}{()}</code>
$E_6^1$		<code>\dynkin[extended]{E}{6}</code>
$E_7^1$		<code>\dynkin[extended]{E}{7}</code>
$E_8^1$		<code>\dynkin[extended]{E}{8}</code>
$F_4^1$		<code>\dynkin[extended]{F}{4}</code>
$G_2^1$		<code>\dynkin[extended]{G}{2}</code>

## 13. AFFINE TWISTED AND UNTWISTED DYNKIN DIAGRAMS

The affine Dynkin diagrams are described in the notation of Kac [12] p. 55:

## Affine Dynkin diagrams

```
\(A^{(1)}_7=\dynkin{A}[1]{7}, \
E^{(2)}_6=\dynkin{E}[2]{6}, \
D^{(3)}_4=\dynkin{D}[3]{4})\)
```

$$A_7^{(1)} = \text{Diagram of } A_7^{(1)}, E_6^{(2)} = \text{Diagram of } E_6^{(2)}, D_4^{(3)} = \text{Diagram of } D_4^{(3)}$$

Table 9: The affine Dynkin diagrams





$A_1^1$		<code>\dynkin{A}[1]{1}</code>
$A_n^1$		<code>\dynkin{A}[1]{}</code>
$B_n^1$		<code>\dynkin{B}[1]{}</code>
$C_n^1$		<code>\dynkin{C}[1]{}</code>
$D_n^1$		<code>\dynkin{D}[1]{}</code>
$E_6^1$		<code>\dynkin{E}[1]{6}</code>
$E_7^1$		<code>\dynkin{E}[1]{7}</code>
$E_8^1$		<code>\dynkin{E}[1]{8}</code>
$F_4^1$		<code>\dynkin{F}[1]{4}</code>
$G_2^1$		<code>\dynkin{G}[1]{2}</code>
$A_2^2$		<code>\dynkin{A}[2]{2}</code>
$A_{ev}^2$		<code>\dynkin{A}[2]{even}</code>
$A_{od}^2$		<code>\dynkin{A}[2]{odd}</code>
$D_n^2$		<code>\dynkin{D}[2]{}</code>
$E_6^2$		<code>\dynkin{E}[2]{6}</code>
$D_4^3$		<code>\dynkin{D}[3]{4}</code>

Table 10: Some more affine Dynkin diagrams

$A_4^2$		<code>\dynkin{A}[2]{4}</code>
---------	---	-------------------------------

continued ...

Table 10: ... continued

$A_5^2$		<code>\dynkin{A}{2}{5}</code>
$A_6^2$		<code>\dynkin{A}{2}{6}</code>
$A_7^2$		<code>\dynkin{A}{2}{7}</code>
$A_8^2$		<code>\dynkin{A}{2}{8}</code>
$D_3^2$		<code>\dynkin{D}{2}{3}</code>
$D_4^2$		<code>\dynkin{D}{2}{4}</code>
$D_5^2$		<code>\dynkin{D}{2}{5}</code>
$D_6^2$		<code>\dynkin{D}{2}{6}</code>
$D_7^2$		<code>\dynkin{D}{2}{7}</code>
$D_8^2$		<code>\dynkin{D}{2}{8}</code>
$D_4^3$		<code>\dynkin{D}{3}{4}</code>
$E_6^2$		<code>\dynkin{E}{2}{6}</code>

## 14. EXTENDED COXETER DIAGRAMS

Extended and Coxeter options together



`\dynkin[extended,Coxeter]{F}{4}`

Table 11: The extended (affine) Coxeter diagrams

$A_n$		<code>\dynkin[extended,Coxeter]{A}{}</code>
$B_n$		<code>\dynkin[extended,Coxeter]{B}{}</code>
$C_n$		<code>\dynkin[extended,Coxeter]{C}{}</code>
$D_n$		<code>\dynkin[extended,Coxeter]{D}{}</code>
$E_6$		<code>\dynkin[extended,Coxeter]{E}{6}</code>
$E_7$		<code>\dynkin[extended,Coxeter]{E}{7}</code>

continued ...

Table 11: ... continued

$E_8$		<code>\dynkin[extended,Coxeter]{E}{8}</code>
$F_4$		<code>\dynkin[extended,Coxeter]{F}{4}</code>
$G_2$		<code>\dynkin[extended,Coxeter]{G}{2}</code>
$H_3$		<code>\dynkin[extended,Coxeter]{H}{3}</code>
$H_4$		<code>\dynkin[extended,Coxeter]{H}{4}</code>
$I_1$		<code>\dynkin[extended,Coxeter]{I}{1}</code>

## 15. KAC STYLE

We include a style called `Kac` which tries to imitate the style of [12].

Kac style

```
\dynkin[Kac]{F}{4}
```









Table 12: The Dynkin diagrams of the extended simple root systems in Kac style. At the moment, it only works on a white background.

$A_1^1$		<code>\dynkin[extended]{A}{1}</code>
$A_n^1$		<code>\dynkin[extended]{A}{}{}</code>
$B_n^1$		<code>\dynkin[extended]{B}{}{}</code>
$C_n^1$		<code>\dynkin[extended]{C}{}{}</code>
$D_n^1$		<code>\dynkin[extended]{D}{}{}</code>
$E_6^1$		<code>\dynkin[extended]{E}{6}</code>
$E_7^1$		<code>\dynkin[extended]{E}{7}</code>
$E_8^1$		<code>\dynkin[extended]{E}{8}</code>

continued ...

Table 12: ... continued

$$\begin{array}{ll} F_4^1 & \circ\text{---}\circ\text{---}\circ\Rightarrow\circ\text{---}\circ \\ G_2^1 & \circ\text{---}\circ\Rightarrow\circ \end{array}$$


```
\dynkin[extended]{F}{4}
\dynkin[extended]{G}{2}
```

## 16. FOLDED DYNKIN DIAGRAMS

The Dynkin diagrams package has limited support for folding Dynkin diagrams.


## Folding

```
\dynkin[fold]{A}{13}
```




## Big fold radius

```
\dynkin[fold,foldradius=1cm]{A}{13}
```



## Small fold radius


```
\dynkin[fold,foldradius=.2cm]{A}{13}
```



Some Dynkin diagrams have multiple foldings, which we attempt to distinguish (not entirely successfully) by their *ply*: the maximum number of roots folded together. Most diagrams can only allow a 2-ply folding, so `fold` is a synonym for `ply=2`.


## 3-ply

```
\dynkin[ply=3]{D}{4}
\dynkin[ply=3]{D}[1]{4}
```



4-ply


```
\dynkin[ply=4]{D}[1]{4}
```



The  $D_\ell^{(1)}$  diagrams can be folded on their left end and separately on their right end:

Left, right and both

```
\dynkin{D}[1]{} \
\dynkin[foldleft]{D}[1]{} \
\dynkin[foldright]{D}[1]{} \
\dynkin[fold]{D}[1]{}
```



We have to be careful about the 4-ply foldings of  $D_{2\ell}^{(1)}$ , for which we can have two different patterns, so by default, the package only draws as much as it can without distinguishing the two:

Default  $D_{2\ell}^{(1)}$  and the two ways to finish it

```
\begin{tikzpicture}
\dynkin[ply=4]{D}[1]{****.*****.*****}%
\end{tikzpicture} \
\begin{tikzpicture}
\dynkin[ply=4]{D}[1]{****.*****.*****}%
\dynkinFold[bend right=65]{1}{13}%
\dynkinFold[bend right=65]{0}{14}%
\end{tikzpicture} \
\begin{tikzpicture}
\dynkin[ply=4]{D}[1]{****.*****.*****}%
\dynkinFold{0}{1}%
\dynkinFold{1}{13}%
\dynkinFold{13}{14}%
\end{tikzpicture}
```





Table 13: Some foldings of Dynkin diagrams

$A_3$		<code>\dynkin[fold]{A}{0}{3}</code>
$C_2$		<code>\dynkin{C}{0}{2}</code>
$A_{2\ell}$		<code>\dynkin[fold]{A}{**.*****.**}</code>
$C_\ell$		<code>\dynkin{C}{}</code>
$B_3$		<code>\dynkin[fold]{B}{0}{3}</code>
$G_2$		<code>\dynkin[reverseArrows]{G}{0}{2}</code>
$D_4$		<code>\dynkin[ply=3]{D}{4}</code>
$G_2$		<code>\dynkin{G}{2}</code>
$D_{\ell+1}$		<code>\dynkin[fold]{D}{}</code>
$B_\ell$		<code>\dynkin{B}{}</code>
$E_6$		<code>\dynkin[fold]{E}{0}{6}</code>
$F_4$		<code>\dynkin[reverseArrows]{F}{0}{4}</code>
$A_3^1$		<code>\dynkin[ply=4]{A}{1}{3}</code>
$A_1^1$		<code>\dynkin{A}{1}{1}</code>
$A_{2\ell}^1$		<code>\dynkin[fold]{A}{1}{**.*****.**}</code>
$C_\ell^1$		<code>\dynkin{C}{1}{}</code>




continued ...

Table 13: ...continued

$B_3^1$		<code>\dynkin[ply=3]{B}{1}{3}</code>
$A_2^2$		<code>\dynkin{A}{2}{2}</code>
$B_3^1$		<code>\dynkin[ply=2]{B}{1}{3}</code>
$G_2^1$		<code>\dynkin{G}{1}{2}</code>
$B_\ell^1$		<code>\dynkin[fold]{B}{1}{}</code>
$D_\ell^2$		<code>\dynkin{D}{2}{}</code>
$D_4^1$		<code>\dynkin[ply=3]{D}{1}{4}</code>
$B_3^1$		<code>\dynkin{B}{1}{3}</code>
$D_4^1$		<code>\dynkin[ply=3]{D}{1}{4}</code>
$G_2^1$		<code>\dynkin{G}{1}{2}</code>
$D_{\ell+}^1$		<code>\dynkin[fold]{D}{1}{}</code>
$D_\ell^2$		<code>\dynkin{D}{2}{}</code>
$D_{\ell+}^1$		<code>\dynkin[foldright]{D}{1}{}</code>
$B_\ell^1$		<code>\dynkin{B}{1}{}</code>

continued ...

Table 13: ...continued

$D_{2\ell}^1$		\begin{tikzpicture}[baseline=0pt] \dynkin[ply=4]{D}{1}{****.*****.*****} \dynkinFold{0}{1} \dynkinFold{1}{13} \dynkinFold{13}{14} \end{tikzpicture}
$A_{\text{od}}^2$		\dynkin{A}{2}{odd}
$D_{2\ell}^1$		\begin{tikzpicture}[baseline=0pt] \dynkin[ply=4]{D}{1}{****.*****.*****} \dynkinFold[bend right=65]{1}{13} \dynkinFold[bend right=65]{0}{14} \end{tikzpicture}
$A_{\text{ev}}^2$		\dynkin{A}{2}{even}
$E_6^1$		\dynkin{fold}{E}{1}{6}
$F_4^1$		\dynkin{reverseArrows}{F}{1}{4}
$E_6^1$		\dynkin{ply=3}{E}{1}{6}
$D_4^3$		\dynkin{D}{3}{4}
$E_7^1$		\dynkin{fold}{E}{1}{7}
$E_6^2$		\dynkin{E}{2}{6}
$F_4^1$		\dynkin{fold}{F}{1}{4}
$G_2^1$		\dynkin{G}{1}{2}

continued ...


Table 13: ...continued

$A_{\text{od}}^2$		<code>\dynkin[odd,fold]{A}{2}{****.***}</code>
$A_{\text{ev}}^2$		<code>\dynkin{A}{2}{even}</code>
$D_3^2$		<code>\dynkin[fold]{D}{2}{3}</code>
$A_2^2$		<code>\dynkin{A}{2}{2}</code>













## 17. ROOT ORDERING

Root ordering

```
\dynkin[label,ordering=Adams]{E}{6}
\dynkin[label,ordering=Bourbaki]{E}{6}
\dynkin[label,ordering=Carter]{E}{6}
\dynkin[label,ordering=Dynkin]{E}{6}
\dynkin[label,ordering=Kac]{E}{6}
```



Default is Bourbaki.

	Adams	Bourbaki	Carter	Dynkin	Kac
$E_6$					
$E_7$					
$E_8$					
$F_4$					
$G_2$					

## 18. CONNECTING DYNKIN DIAGRAMS

We can make some sophisticated folded diagrams by drawing multiple diagrams, each with a name:

Name a diagram

```
\dynkin[name=Bob]{D}{6}
```



We can then connect the two with folding edges:


Connect diagrams

```
\begin{tikzpicture}
    \dynkin[name=upper]{A}{3}
    \node (current) at ($(upper root 1)+(0,-.3cm)$) {};
    \dynkin[at=(current),name=lower]{A}{3}
    \begin{scope}[on background layer]
        \foreach \i in {1,...,3}%
        {
            \draw[/Dynkin diagram/foldStyle]
                ($(upper root \i)$) -- ($(lower
root \i)$);
        }
    \end{scope}
\end{tikzpicture}
```




The following diagrams arise in the Satake diagrams of the pseudo-Riemannian symmetric spaces [1].

```
\pgfkeys{/Dynkin diagram,edgeLength=.5cm,foldradius=.5cm}
\begin{tikzpicture}
    \dynkin[name=1]{A}{IIIb}
    \node (a) at (.3,.4){};
    \dynkin[name=2,at=(a)]{A}{IIIb}
    \begin{scope}[on background layer]
        \foreach \i in {1,...,7}%
        {
            \draw[/Dynkin diagram/foldStyle]
                ($(1 root \i)$)
                --
                ($(2 root \i)$);
        }
    \end{scope}
\end{tikzpicture}
```



```
\pgfkeys{/Dynkin diagram/edgeLength=.75cm,/Dynkin
    diagram/edge/.style={draw=white,double=black,very thick},
}
\begin{tikzpicture}
    \foreach \d in {1,\dots,4}
    {
        \node (current) at ($(\d*.05,\d*.3)$){};
        \dynkin[name=\d,at=(current)]{D}{oo..ooo}
    }
    \begin{scope}[on background layer]
        \foreach \i in {1,\dots,6}%
        {%
            \draw[/Dynkin diagram/foldStyle] ($(1 root
\i)$) -- ($(2 root \i)$);%
            \draw[/Dynkin diagram/foldStyle] ($(2 root
\i)$) -- ($(3 root \i)$);%
            \draw[/Dynkin diagram/foldStyle] ($(3 root
\i)$) -- ($(4 root \i)$);%
        }%
    \end{scope}
\end{tikzpicture}
```




## 19. OTHER EXAMPLES


Below we draw the Vogan diagrams of some affine Lie superalgebras [17, 16].

$\mathfrak{sl}(2m|2n)^{(2)}$ 


```
\begin{tikzpicture}
\dynkin[ply=2,label]{B}{1}{oo.oto.oo}
\dynkinLabelRoot*{7}{1}
\end{tikzpicture}
```




```
\dynkin[label]{B}{1}{oo.oto.oo}
```




```
\dynkin[ply=2,label]{B}{1}{oo.Oto.Oo}
```




```
\dynkin[label]{B}{1}{oo.Oto.Oo}
```




```
\dynkin[label]{D}{1}{oo.oto.ooo}
```



```
\dynkin[label]{D}{1}{oo.ot0.ooo}
```




```
\dynkin[label,fold]{D}{1}{oo.oto.ooo}
```




$\mathfrak{sl}(2m+1|2n)^2$


```
\dynkin[label]{B}{1}{oo.oto.oo}
```



```
\dynkin[label]{B}{1}{o0.oto.oo}
```




```
\dynkin[label,fold]{B}{1}{oo.oto.oo}
```




$\mathfrak{sl}(2m+1|2n+1)^2$

```
\dynkin[label]{D}{2}{o.oto.oo}
```




```
\dynkin[label]{D}{2}{o.0t0.oo}
```




$\mathfrak{sl}(2|2n+1)^{(2)}$ 


```
\dynkin[ply=2,label,doubleEdges]{B}[1]{oo.0to.0o}
```




```
\dynkin[ply=2,label,doubleFold]{B}[1]{oo.0to.0o}
```




```
\dynkin[ply=2,label,doubleEdges]{B}[1]{oo.0t0.oo}
```




```
\dynkin[ply=2,label,doubleFold]{B}[1]{oo.0t0.oo}
```


 $\mathfrak{sl}(2|2n)^{(2)}$ 

```
\dynkin[ply=2,label,doubleEdges]{D}[1]{oo.oto.ooo}
```




```
\dynkin[ply=2,label,doubleFoldLeft]{D}[1]{oo.oto.ooo}
```




$\mathfrak{osp}(2m|2n)^{(2)}$ 


```
\dynkin[label,labelMacro/.code={1}]{D}[2]{o.oto.oo}
```



```
\dynkin[label,labelMacro/.code={1}]{D}[2]{o.0to.0o}
```


 $\mathfrak{osp}(2|2n)^{(2)}$ 


```
\dynkin[label,labelMacro/.code=\labelIt{\#1},
affineMark=*]
{D}[2]{o.o.o.o*}
```




```
\dynkin[label,labelMacro/.code=\labelIt{\#1},
affineMark=*]
{D}[2]{o.0.o.o*}
```


 $\mathfrak{sl}(1|2n+1)^4$ 

```
\dynkin[label,labelMacro/.code={1}]{D}[2]{o.o.o.o*}
```




```
\dynkin[label,labelMacro/.code={1}]{D}[2]{o.o.0.o*}
```




$A^1$ 


```
\begin{tikzpicture}
  \dynkin[name=upper]{A}{oo.t.oo}
  \node (Dynkin current) at (upper root 1){};
  \dynkinSouth
  \dynkin[at=(Dynkin current),name=lower]{A}{oo.t.oo}
  \begin{scope}[on background layer]
  \foreach \i in {1,...,5}{
    \draw[/Dynkin diagram/foldStyle]
    ($(\text{upper root }\i)$) --
    ($(\text{lower root }\i)$);
  }
  \end{scope}
\end{tikzpicture}
```




```
\dynkin[fold]{A}[1]{oo.t.ooooo.t.oo}
```




```
\dynkin[fold,affineMark=t]{A}[1]{oo.o.oootoo.o.oo}
```




```
\dynkin[affineMark=t]{A}[1]{o*.t.*o}
```

 $B^1$ 


```
\dynkin[affineMark=*]{A}[2]{o.oto.o*}
```



```
\dynkin[affineMark=*]{A}[2]{o.oto.o*}
```




```
\dynkin[affineMark=*]{A}[2]{o.ooo.oo}
```




```
\dynkin[odd]{A}[2]{oo.*to.*o}
```



```
\dynkin[odd,fold]{A}[2]{oo.oto.oo}
```



```
\dynkin[odd,fold]{A}[2]{o*.oto.o*}
```



$D^1$

```
\dynkin{D}{otoo}
```




```
\dynkin{D}{ot*o}
```




```
\dynkin [fold]{D}{otoo}
```

 $C^1$ 


```
\dynkin [doubleEdges,fold,affineMark=t,odd]{A}[2]{to . o*}
```



```
\dynkin [doubleEdges,fold,affineMark=t,odd]{A}[2]{t*.oo}
```


 $F^1$ 

```
\begin{tikzpicture}%
  \dynkin{A}{otoo*}%
  \dynkinQuadrupleEdge{1}{2}%
  \dynkinTripleEdge{4}{3}%
\end{tikzpicture}%
```



```
\begin{tikzpicture}%
  \dynkin{A}{*too}%
  \dynkinQuadrupleEdge{1}{2}%
  \dynkinTripleEdge{4}{3}%
\end{tikzpicture}%
```





$\mathfrak{g}$	diagram	$V$	$\Delta$	$\alpha_i$
$A_n$		$\mathbb{R}^{n+1}/\langle \sum e_j \rangle$	$e_i - e_j$	$e_i - e_{i+1}$
$B_n$		$\mathbb{R}^n$	$\pm e_i, \pm e_i \pm e_j, i \neq j$	$e_i - e_{i+1}, e_n$
$C_n$		$\mathbb{R}^n$	$\pm 2e_i, \pm e_i \pm e_j, i \neq j$	$e_i - e_{i+1}, 2e_n$
$D_n$		$\mathbb{R}^n$	$\pm e_i \pm e_j, i \neq j$	$\begin{cases} e_i - e_{i+1}, & i \leq n-2 \\ e_{n-1} + e_n & \end{cases}$

$\mathfrak{g}$	diagram	$V$	$\Delta$	$\alpha_i$
$E_8$		$\mathbb{R}^8$	$\begin{cases} \pm 2e_i \pm 2e_j, & i \neq j, \\ \sum_i (-1)^{m_i} e_i, & \sum m_i \text{ even} \end{cases}$	$\begin{cases} 2e_1 - 2e_2, \\ 2e_2 - 2e_3, \\ 2e_3 - 2e_4, \\ 2e_4 - 2e_5, \\ 2e_5 - 2e_6, \\ 2e_6 + 2e_7, \\ -\sum e_j, \\ 2e_6 - 2e_7 \end{cases}$
$E_7$		$\mathbb{R}^8 / \langle e_1 - e_2 \rangle$	quotient of $E_8$	quotient of $E_8$
$E_6$		$\mathbb{R}^8 / \langle e_1 - e_2, e_2 - e_3 \rangle$	quotient of $E_8$	quotient of $E_8$
$F_4$		$\mathbb{R}^4$	$\begin{cases} \pm 2e_i, \\ \pm 2e_i \pm 2e_j, i \neq j, \\ \pm e_1 \pm e_2 \pm e_3 \pm e_4 \end{cases}$	$\begin{cases} 2e_2 - 2e_3, \\ 2e_3 - 2e_4, \\ 2e_4, \\ e_1 - e_2 - e_3 - e_4 \end{cases}$
$G_2$		$\mathbb{R}^3 / \langle \sum e_j \rangle$	$\begin{cases} \pm(1, -1, 0), \\ \pm(-1, 0, 1), \\ \pm(0, -1, 1), \\ \pm(2, -1, -1), \\ \pm(1, -2, 1), \\ \pm(-1, -1, 2) \end{cases}$	$\begin{cases} (-1, 0, 1), \\ (2, -1, -1) \end{cases}$

## 20. SYNTAX

The syntax is `\dynkin[<options>]{<letter>}[<twisted rank>]{<rank>}` where `<letter>` is A, B, C, D, E, F or G, the family of root system for the Dynkin diagram, `<twisted rank>` is 0, 1, 2, 3 (default is 0) representing:

- 0 finite root system
- 1 affine extended root system, i.e. of type <sup>(1)</sup>
- 2 affine twisted root system of type <sup>(2)</sup>
- 3 affine twisted root system of type <sup>(3)</sup>

and `<rank>` is

- (1) an integer representing the rank or
- (2) blank to represent an indefinite rank or
- (3) the name of a Satake diagram as in section 4.

## 21. OPTIONS

```
text/.style = {TikZ style data},
default : scale=.7
      Style for any labels on the roots.
name = <string>,
default : anonymous
```

continued ...

Table 16: ... continued

A name for the Dynkin diagram, with `anonymous` treated as a blank; see section 18.

`parabolic = <integer>,`  
`default : 0`  
     A parabolic subgroup with specified integer, where the integer is computed as  $n = \sum 2^{i-1}a_i$ ,  $a_i = 0$  or  $1$ , to say that root  $i$  is crossed, i.e. a noncompact root.

`radius = <number>cm,`  
`default : .05cm`  
     size of the dots and of the crosses in the Dynkin diagram

`edgeLength = <number>cm,`  
`default : .35cm`  
     distance between nodes in the Dynkin diagram

`edge/.style = TikZ style data,`  
`default : thin`  
     style of edges in the Dynkin diagram

`mark = <o,O,t,x,X,*>,`  
`default : *`  
     default root mark

`affineMark = o,O,t,x,X,*,`  
`default : *`  
     default root mark for root zero in an affine Dynkin diagram

`label = true or false,`  
`default : false`  
     whether to label the roots according to the current labelling scheme.

`labelMacro = <1-parameter TeX macro>,`  
`default : #1`  
     the current labelling scheme.

`makeIndefiniteEdge = <edge pair  $i-j$  or list of such>,`  
`default : {}`  
     edge pair or list of edge pairs to treat as having indefinitely many roots on them.

`indefiniteEdgeRatio = <float>,`  
`default : 1.6`  
     ratio of indefinite edge lengths to other edge lengths.

`indefiniteEdge/.style = <TikZ style data>,`  
`default : draw=black,fill=white,thin,densely dotted`  
     style of the dotted or dashed middle third of each indefinite edge.

`arrows = <true or false>,`  
`default : true`  
     whether to draw the arrows that arise along the edges.

`reverseArrows = <true or false>,`  
`default : true`  
     whether to reverse the direction of the arrows that arise along the edges.

`fold = <true or false>,`

continued ...

Table 16: ... continued

```

default : true
        whether, when drawing Dynkin diagrams, to draw them 2-ply.
ply =  $\langle 0, 1, 2, 3, 4 \rangle$ ,
default : 0
        how many roots get folded together, at most.
foldleft =  $\langle \text{true or false} \rangle$ ,
default : true
        whether to fold the roots on the left side of a Dynkin diagram.
foldright =  $\langle \text{true or false} \rangle$ ,
default : true
        whether to fold the roots on the right side of a Dynkin diagram.
foldradius =  $\langle \text{length} \rangle$ ,
default : .3cm
        the radius of circular arcs used in curved edges of folded Dynkin
        diagrams.
foldStyle =  $\langle \text{TikZ style data} \rangle$ ,
default : draw=black!40,fill=none,line width=radius
        when drawing folded diagrams, style for the fold indicators.
 $*/.$ style =  $\langle \text{TikZ style data} \rangle$ ,
default : draw=black,fill=black
        style for roots like •
 $o/.$ style =  $\langle \text{TikZ style data} \rangle$ ,
default : draw=black,fill=black
        style for roots like o
 $0/.$ style =  $\langle \text{TikZ style data} \rangle$ ,
default : draw=black,fill=black
        style for roots like @
 $t/.$ style =  $\langle \text{TikZ style data} \rangle$ ,
default : draw=black,fill=black
        style for roots like *
 $x/.$ style =  $\langle \text{TikZ style data} \rangle$ ,
default : draw=black
        style for roots like x
 $X/.$ style =  $\langle \text{TikZ style data} \rangle$ ,
default : draw=black,thick
        style for roots like x
leftFold/.style =  $\langle \text{TikZ style data} \rangle$ ,
default :
        style to override the fold style when folding roots together on the
        left half of a Dynkin diagram
rightFold/.style =  $\langle \text{TikZ style data} \rangle$ ,
default :
        style to override the fold style when folding roots together on the
        right half of a Dynkin diagram
doubleEdges =  $\langle \rangle$ ,
default : not set

```

continued ...

Table 16: ... continued

<code>doubleFold = ⟨⟩,</code>	set to override the <code>fold</code> style when folding roots together in a Dynkin diagram, so that the foldings are indicated with double edges (like those of an $F_4$ Dynkin diagram without arrows).
<code>default : not set</code>	
	set to override the <code>fold</code> style when folding roots together in a Dynkin diagram, so that the foldings are indicated with double edges (like those of an $F_4$ Dynkin diagram without arrows), but filled in solidly.
<code>doubleLeft = ⟨⟩,</code>	
<code>default : not set</code>	set to override the <code>fold</code> style when folding roots together at the left side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an $F_4$ Dynkin diagram without arrows).
<code>doubleFoldLeft = ⟨⟩,</code>	
<code>default : not set</code>	set to override the <code>fold</code> style when folding roots together at the left side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an $F_4$ Dynkin diagram without arrows), but filled in solidly.
<code>doubleRight = ⟨⟩,</code>	
<code>default : not set</code>	set to override the <code>fold</code> style when folding roots together at the right side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an $F_4$ Dynkin diagram without arrows).
<code>doubleFoldRight = ⟨⟩,</code>	
<code>default : not set</code>	set to override the <code>fold</code> style when folding roots together at the right side of a Dynkin diagram, so that the foldings are indicated with double edges (like those of an $F_4$ Dynkin diagram without arrows), but filled in solidly.
<code>Coxeter = ⟨true or false⟩,</code>	
<code>default : false</code>	whether to draw a Coxeter diagram, rather than a Dynkin diagram.
<code>ordering = ⟨Adams, Bourbaki, Carter, Dynkin, Kac⟩,</code>	
<code>default : Bourbaki</code>	which ordering of the roots to use in exceptional root systems as in section 17.

All other options are passed to TikZ.

## REFERENCES

1. Kurando Baba, *Satake diagrams and restricted root systems of semisimple pseudo-Riemannian symmetric spaces*, Tokyo J. Math. **32** (2009), no. 1, 127–158. MR 2541161

2. Nicolas Bourbaki, *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressley. MR 1890629
3. R. W. Carter, *Lie algebras of finite and affine type*, Cambridge Studies in Advanced Mathematics, vol. 96, Cambridge University Press, Cambridge, 2005. MR 2188930
4. Meng-Kiat Chuah, *Cartan automorphisms and Vogan superdiagrams*, Math. Z. **273** (2013), no. 3–4, 793–800. MR 3030677
5. Cristina Draper Fontanals and Valerio Guido, *On the real forms of the exceptional Lie algebra  $\mathfrak{e}_6$  and their Satake diagrams*, Non-associative and non-commutative algebra and operator theory, Springer Proc. Math. Stat., vol. 160, Springer, Cham, 2016, pp. 211–226. MR 3613831
6. E. B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, Mat. Sbornik N.S. **30(72)** (1952), 349–462 (3 plates), Reprinted in English translation in [7]. MR 0047629
7. ———, *Selected papers of E. B. Dynkin with commentary*, American Mathematical Society, Providence, RI; International Press, Cambridge, MA, 2000, Edited by A. A. Yushkevich, G. M. Seitz and A. L. Onishchik. MR 1757976
8. L. Frappat, A. Sciarrino, and P. Sorba, *Structure of basic Lie superalgebras and of their affine extensions*, Comm. Math. Phys. **121** (1989), no. 3, 457–500. MR 990776
9. L. C. Grove and C. T. Benson, *Finite reflection groups*, second ed., Graduate Texts in Mathematics, vol. 99, Springer-Verlag, New York, 1985. MR 777684
10. Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics, vol. 34, American Mathematical Society, Providence, RI, 2001, Corrected reprint of the 1978 original. MR 1834454
11. James E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, vol. 29, Cambridge University Press, Cambridge, 1990. MR 1066460
12. Victor G. Kac, *Infinite-dimensional Lie algebras*, third ed., Cambridge University Press, Cambridge, 1990. MR 1104219
13. S. Pratik Khastgir and Ryu Sasaki, *Non-canonical folding of Dynkin diagrams and reduction of affine Toda theories*, Progr. Theoret. Phys. **95** (1996), no. 3, 503–518. MR 1388245
14. A. L. Onishchik and È. B. Vinberg, *Lie groups and algebraic groups*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1990, Translated from the Russian and with a preface by D. A. Leites. MR 91g:22001
15. A. L. Onishchik and È. B. Vinberg, *Lie groups and algebraic groups*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1990, Translated from the Russian and with a preface by D. A. Leites. MR 1064110
16. B. Ransingh, *Vogan diagrams of affine twisted Lie superalgebras*, ArXiv e-prints (2013).
17. Biswajit Ransingh, *Vogan diagrams of untwisted affine Kac-Moody superalgebras*, Asian-Eur. J. Math. **6** (2013), no. 4, 1350062, 10. MR 3149279
18. V. Regelskis and B. Vlaar, *Reflection matrices, coideal subalgebras and generalized Satake diagrams of affine type*, ArXiv e-prints (2016).
19. Ichirô Satake, *Algebraic structures of symmetric domains*, Kanô Memorial Lectures, vol. 4, Iwanami Shoten, Tokyo; Princeton University Press, Princeton, N.J., 1980. MR 591460
20. T. A. Springer, *Linear algebraic groups*, second ed., Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2009. MR 2458469
21. È. B. Vinberg (ed.), *Lie groups and Lie algebras, III*, Encyclopaedia of Mathematical Sciences, vol. 41, Springer-Verlag, Berlin, 1994, Structure of Lie groups and Lie algebras, A translation of it Current problems in mathematics. Fundamental directions. Vol. 41 (Russian), Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1990 [MR1056485 (91b:22001)], Translation by V. Minachin [V. V. Minakhin], Translation edited by A. L. Onishchik and È. B. Vinberg. MR 1349140
22. Jean-Bernard Zuber, *Generalized Dynkin diagrams and root systems and their folding*, Topological field theory, primitive forms and related topics (Kyoto, 1996), Progr. Math., vol. 160, Birkhäuser Boston, Boston, MA, 1998, pp. 453–493. MR 1653035