

Sample Document Using Interchangable Numbering

Nicola Talbot

July 31, 2014

Abstract

This is a sample document illustrating the use of the `glossaries` package. The functions here have been taken from “Tables of Integrals, Series, and Products” by I.S. Gradshteyn and I.M. Ryzhik.

The glossary lists both page numbers and equation numbers. Since the majority of the entries use the equation number, `counter=equation` was used as a package option. Note that this example will only work where the page number and equation number compositor is the same. So it won’t work if, say, the page numbers are of the form 2-4 and the equation numbers are of the form 4.6. As most of the glossary entries should have an italic format, it is easiest to set the default format to italic.

Contents

Special Functions	2
1 Gamma Functions	4
2 Error Functions	6
3 Beta Function	7
4 Chebyshev's polynomials	8
5 Hermite polynomials	9
6 Laguerre polynomials	10
7 Bessel Functions	11
8 Confluent hypergeometric function	12
9 Parabolic cylinder functions	13
10 Elliptical Integral of the First Kind	14
11 Constants	15

Index of Special Functions and Notations

Numbers in *italic* indicate the equation number, numbers in **bold** indicate page numbers where the main definition occurs.

Notation	Function Name	
$B(x, y)$	Beta function	<i>3.1–3.3</i>
$B_x(p, q)$	Incomplete beta function	<i>3.4</i>
C	Euler’s constant	<i>11.1</i>
$D_p(z)$	Parabolic cylinder functions	<i>9.1</i>
$\operatorname{erf}(x)$	Error function	<i>2.1</i> , 6
$\operatorname{erfc}(x)$	Complementary error function	<i>2.2</i>
$F(\phi, k)$	Elliptical integral of the first kind	<i>10.1</i>
G	Catalan’s constant	<i>11.2</i>
$\Gamma(z)$	Gamma function	<i>1.1</i> , <i>1.2</i> , <i>1.5</i> , 4
$\Gamma(\alpha, x)$	Incomplete gamma function	<i>1.4</i>
$\gamma(\alpha, x)$	Incomplete gamma function	<i>1.3</i>
$H_n(x)$	Hermite polynomials	<i>5.1</i>
$k_\nu(x)$	Bateman’s function	<i>8.2</i>
$L_n^\alpha(x)$	Laguerre polynomials	<i>6.1</i>

Notation	Function Name	
$\Phi(\alpha, \gamma; z)$	confluent hypergeometric function	<i>8.1</i>
$\psi(x)$	Psi function	<i>1.6</i>
$T_n(x)$	Chebyshev's polynomials of the first kind	<i>4.1</i>
$U_n(x)$	Chebyshev's polynomials of the second kind	<i>4.2</i>
$Z_\nu(z)$	Bessel functions	<i>7.1</i>

Chapter 1

Gamma Functions

The **gamma function** is defined as

$$\Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt \quad (1.1)$$

$$\Gamma(x+1) = x\Gamma(x) \quad (1.2)$$

$$\gamma(\alpha, x) = \int_0^x e^{-t} t^{\alpha-1} dt \quad (1.3)$$

$$\Gamma(\alpha, x) = \int_x^\infty e^{-t} t^{\alpha-1} dt \quad (1.4)$$

$$\Gamma(\alpha) = \Gamma(\alpha, x) + \gamma(\alpha, x) \quad (1.5)$$

$$\psi(x) = \frac{d}{dx} \ln \Gamma(x) \quad (1.6)$$

Chapter 2

Error Functions

The **error function** is defined as:

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt \quad (2.1)$$

$$\text{erfc}(x) = 1 - \text{erf}(x) \quad (2.2)$$

Chapter 3

Beta Function

$$B(x, y) = 2 \int_0^1 t^{x-1} (1-t^2)^{y-1} dt \quad (3.1)$$

Alternatively:

$$B(x, y) = 2 \int_0^{\frac{\pi}{2}} \sin^{2x-1} \phi \cos^{2y-1} \phi d\phi \quad (3.2)$$

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = B(y, x) \quad (3.3)$$

$$B_x(p, q) = \int_0^x t^{p-1} (1-t)^{q-1} dt \quad (3.4)$$

Chapter 4

Chebyshev's polynomials

$$T_n(x) = \cos(n \arccos x) \tag{4.1}$$

$$U_n(x) = \frac{\sin[(n+1) \arccos x]}{\sin[\arccos x]} \tag{4.2}$$

Chapter 5

Hermite polynomials

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2}) \quad (5.1)$$

Chapter 6

Laguerre polynomials

$$L_n^\alpha(x) = \frac{1}{n!} e^x x^{-\alpha} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}) \quad (6.1)$$

Chapter 7

Bessel Functions

Bessel functions $Z_\nu(z)$ are solutions of

$$\frac{d^2 \textcolor{red}{Z}_\nu}{dz^2} + \frac{1}{z} \frac{dZ_\nu}{dz} + \left(1 - \frac{\nu^2}{z^2} Z_\nu = 0\right) \quad (7.1)$$

Chapter 8

Confluent hypergeometric function

$$\Phi(\alpha, \gamma; z) = 1 + \frac{\alpha}{\gamma} \frac{z}{1!} + \frac{\alpha(\alpha+1)}{\gamma(\gamma+1)} \frac{z^2}{2!} + \frac{\alpha(\alpha+1)(\alpha+2)}{\gamma(\gamma+1)(\gamma+2)} \frac{z^3}{3!} + \cdots \quad (8.1)$$

$$k_\nu(x) = \frac{2}{\pi} \int_0^{\pi/2} \cos(x \tan \theta - \nu \theta) d\theta \quad (8.2)$$

Chapter 9

Parabolic cylinder functions

$$D_p(z) = 2^{\frac{p}{2}} e^{-\frac{z^2}{4}} \left\{ \frac{\sqrt{\pi}}{\Gamma\left(\frac{1-p}{2}\right)} \Phi\left(-\frac{p}{2}, \frac{1}{2}; \frac{z^2}{2}\right) - \frac{\sqrt{2\pi}z}{\Gamma\left(-\frac{p}{2}\right)} \Phi\left(\frac{1-p}{2}, \frac{3}{2}; \frac{z^2}{2}\right) \right\} \quad (9.1)$$

Chapter 10

Elliptical Integral of the First Kind

$$F(\phi, k) = \int_0^\phi \frac{d\alpha}{\sqrt{1 - k^2 \sin^2 \alpha}} \quad (10.1)$$

Chapter 11

Constants

$$\textcolor{red}{C} = 0.577\,215\,664\,901\dots \tag{11.1}$$

$$\textcolor{red}{G} = 0.915\,965\,594\dots \tag{11.2}$$